

| Kentucky Core Healthcare Measures Set | | |
|--|-------|----------|
| | | |
| | | |
| Preventive Care | NQF # | Priority |
| Cervical Cancer Screening | 32 | High |
| Colorectal Cancer Screening | 34 | High |
| Breast Cancer Screening | 2372 | High |
| Body Mass Index (BMI) Screening and Follow-Up | 421 | High |
| Influenza Immunization | 41 | Standard |
| Chlamydia Screening in Women | 33 | Standard |
| Pediatric Care | NQF # | Status |
| Childhood Immunization Status (Combination 10) | 38 | High |
| Immunizations for Adolescents (Combination 2) | 1407 | High |
| Tobacco Use and Help with Quitting Among Adolescents | 2803 | High |
| Well-Child visits in the 3-6 Years of Life | 1516 | High |
| Well-Child Visits in the First 15 Months of Life | 1392 | High |
| Follow-Up Care for Children Prescribed Attention-Deficit/Hyperactivity Disorder (ADHD) | | |
| Medication | 108 | Standard |
| Appropriate Testing for Children with Pharyngitis | 2 | Standard |
| Appropriate Treatment for Children with Upper Respiratory Infection (URI) | 69 | Standard |
| Contraceptive Care – Most and Moderately Effective Methods: Ages 15–20 | 2903 | Standard |
| Behavioral Health | NQF # | Status |
| Screening for Clinical Depression and Follow Up Plan | 418 | High |
| Tobacco Use: Screening and Cessation Intervention | 28 | High |
| Antidepressant Medication Management | 105 | High |
| Chronic and Acute Care | NQF # | Status |
| Comprehensive Diabetes Care: Hemoglobin (HbA1c) Poor Control (>9.0%) | 59 | High |
| Medication Adherence for Diabetes Medications | 541 | High |
| Statin Therapy for Patients with Diabetes | | High |
| Statin Therapy for Patients with Cardiovascular Disease | | High |
| Controlling High Blood Pressure (Hypertension) | 18 | High |
| Medication Adherence for Hypertension (RAS antagonists) | 541 | High |
| Medication Management for People with Asthma | 1799 | High |
| Medication Reconciliation Post-Discharge | 97 | High |
| Comprehensive Diabetes Care: Medical Attention for Nephropathy | 62 | Standard |
| Comprehensive Diabetes Care: Foot Exam | 56 | Standard |
| Comprehensive Diabetes Care: Eye Exam (Retinal) Performed | 55 | Standard |
| Avoidance of Antibiotic Treatment in Adults with Acute Bronchitis | 58 | Standard |
| Cost and Utilization | NQF# | Status |
| Plan All-Cause Readmissions | 1768 | High |
| Use of Imaging Studies for Low Back Pain | 52 | Standard |

Future Areas of Development

| Behavioral Health | NQF # |
|---|-------|
| Unhealthy Alcohol Use Screening and Follow-up | NA |
| The group recommends the development or selection of an opioid prescribing-related measure that will help to measure inappropriate prescribing patterns, at the physician level. | |
| Although this initial list is meant to be a starting point for measuring behavioral health care in the primary care setting, the group would like to see measures applying to the care of patients with serious mental illnesses in future core sets. | |
| The group recognizes that there is additional need for behavioral health measures designed for the pediatric setting, beyond "Screening for Clinical Depression and Follow Up Plan," for preventive reasons. | |
| The group recommends future selection of a measure that tracks outcome of depression management, such as achieving remission or progress towards remission. | |
| Preventive Care | NQF # |
| CG-CAHPS | 5 |
| Pneumococcal Vaccination Coverage for Older Adults | |
| Pediatric Care | NQF # |
| CG-CAHPS | 5 |
| Developmental Screening in the First Three Years of Life | 1448 |
| Number of eligible patients with preventive dental services | |
| Screening for Depression and Follow-Up Plan: Ages 12–17 (CDF-CH) | 418 |
| Weight Assessment and Counseling for Nutrition and Physical Activity for Children/Adolescents | |
| Chronic and Acute Care | NQF # |
| There is need for the development of a measure for appropriate screening for lung cancer | |