

Immunizations and Developmental Milestones for Your Child from Birth Through 6 Years Old

Child's Name _____

Birth Date _____

		Birth	1 MONTH	2 MONTHS	4 MONTHS	6 MONTHS
Recommended Immunizations	Hepatitis B	☐ HepB	☐ HepB ¹			☐ HepB
	Rotavirus			☐ RV	☐ RV	☐ RV
	Diphtheria, Tetanus, Pertussis			☐ DTaP	☐ DTaP	☐ DTaP
	<i>Haemophilus influenzae</i> type b			☐ Hib	☐ Hib	☐ Hib
	Pneumococcal			☐ PCV	☐ PCV	☐ PCV
	Inactivated Poliovirus			☐ IPV	☐ IPV	☐ IPV
	Influenza (Flu)					☐ Influenza, first dose ² ☐ second dose
Milestones*	Milestones should be achieved by the end of the age indicated. Talk to your child's doctor about age-appropriate milestones if your child was born prematurely.	☐ Recognizes caregiver's voice ☐ Turns head toward breast or bottle ☐ Communicates through body language, fussing or crying	☐ Starts to smile ☐ Raises head when on tummy ☐ Calms down when rocked, cradled or sung to	☐ Begins to smile at people ☐ Coos, makes gurgling sounds ☐ Begins to follow things with eyes ☐ Can hold head up	☐ Babbles with expression ☐ Likes to play with people ☐ Reaches for toy with one hand ☐ Brings hands to mouth	☐ Knows familiar faces ☐ Responds to own name ☐ Brings things to mouth ☐ Rolls over in both directions
Growth	At each well child visit, enter date, length, weight, and percentile information to keep track of your child's progress.	WEIGHT / PERCENTILE	WEIGHT / PERCENTILE	WEIGHT / PERCENTILE	WEIGHT / PERCENTILE	WEIGHT / PERCENTILE
		LENGTH / PERCENTILE	LENGTH / PERCENTILE	LENGTH / PERCENTILE	LENGTH / PERCENTILE	LENGTH / PERCENTILE
		HEAD CIRCUMFERENCE	HEAD CIRCUMFERENCE	HEAD CIRCUMFERENCE	HEAD CIRCUMFERENCE	HEAD CIRCUMFERENCE

continues on back page

 Shaded boxes indicate the vaccine can be given during shown age range.

VISIT DATE

VISIT DATE

VISIT DATE

VISIT DATE

VISIT DATE

¹ The second dose of HepB may be given either at the 1 month or 2 month visit.

² Two doses given at least four weeks apart are recommended for children aged 6 months through 8 years of age who are getting a flu vaccine for the first time and for some other children in this age group.

* Milestones adapted from Bright Futures: Guidelines for Health Supervision of Infants, Children, and Adolescents Third Edition, edited by Joseph Hagan, Jr., Judith S. Shaw, and Paula M. Duncan, 2008, Elk Grove Village, IL: American Academy of Pediatrics.

If your child has any medical conditions that put him at risk for infections or is traveling outside the United States, talk to your child's doctor about additional vaccines that he may need.



U.S. Department of
Health and Human Services
Centers for Disease
Control and Prevention



American Academy
of Pediatrics



DEDICATED TO THE HEALTH OF ALL CHILDREN™

Immunizations and Developmental Milestones for Your Child from Birth Through 6 Years Old

Child's Name _____

Birth Date _____

		12 MONTHS	15 MONTHS	18 MONTHS	19–23 MONTHS	2–3 YEARS	4–6 YEARS
Recommended Immunizations	Hepatitis B	☐ HepB (Final dose administered between 6 and 18 months)					
	Diphtheria, Tetanus, Pertussis		☐ DTaP				☐ DTaP
	Haemophilus influenzae type b	☐ Hib					
	Pneumococcal	☐ PCV					
	Inactivated Poliovirus	☐ IPV					☐ IPV
	Influenza (Flu)	☐ Influenza , first dose ² ☐ second dose (if needed)			Age 2 3 ☐ ☐ Influenza , first dose ² ☐ ☐ second dose (if needed)		Age 4 5 6 ☐ ☐ ☐ Influenza , first dose ² ☐ ☐ ☐ second dose (if needed)
	Measles, Mumps, Rubella	☐ MMR					☐ MMR
	Varicella	☐ Varicella					☐ Varicella
	Hepatitis A	☐ ☐ Hep A ³					
Milestones*	Milestones should be achieved by the end of the age indicated. Talk to your child's doctor about age-appropriate milestones if your child was born prematurely.	☐ Cries when mom or dad leaves ☐ Says "mama" and "dada" ☐ Copies gestures (for example, waves "bye bye") ☐ May stand alone	☐ Imitates what you are doing ☐ Drinks from a cup ☐ Scribbles on his own ☐ Walks well	☐ Points to show others something interesting ☐ Says several single words ☐ Points to one body part ☐ May walk up steps and run	☐ Plays mainly beside other children ☐ Follows two-step commands ☐ Plays simple make-believe games ☐ Throws ball overhand	☐ Can name most familiar things ☐ Shows affection for friends without prompting ☐ Turns book pages one at a time ☐ Kicks a ball	☐ Speaks very clearly ☐ Tells stories ☐ Can print some letters or numbers ☐ Hops; may be able to skip
Growth	At each well child visit, enter date, length, weight, and percentile information to keep track of your child's progress.	WEIGHT / PERCENTILE	WEIGHT / PERCENTILE	WEIGHT / PERCENTILE	WEIGHT / PERCENTILE	WEIGHT	WEIGHT
		LENGTH / PERCENTILE	LENGTH / PERCENTILE	LENGTH / PERCENTILE	LENGTH / PERCENTILE	HEIGHT	HEIGHT
		HEAD CIRCUMFERENCE	HEAD CIRCUMFERENCE	HEAD CIRCUMFERENCE	HEAD CIRCUMFERENCE	BMI	BMI

 Shaded boxes indicate the vaccine can be given during shown age range.

VISIT DATE

VISIT DATE

VISIT DATE

VISIT DATE

VISIT DATE

VISIT DATE

² Two doses given at least four weeks apart are recommended for children aged 6 months through 8 years of age who are getting a flu vaccine for the first time and for some other children in this age group.

³ Two doses of HepA vaccine are needed for lasting protection. The first dose of HepA vaccine should be given between 12 months and 23 months of age. The second dose should be given 6 to 18 months later. HepA vaccination may be given to any child 12 months and older to protect against HepA. Children and adolescents who did not receive the HepA vaccine and are at high-risk, should be vaccinated against HepA.

* Milestones adapted from *Bright Futures: Guidelines for Health Supervision of Infants, Children, and Adolescents* Third Edition, edited by Joseph Hagan, Jr., Judith S. Shaw, and Paula M. Duncan, 2008, Elk Grove Village, IL: American Academy of Pediatrics.

If your child has any medical conditions that put him at risk for infections or is traveling outside the United States, talk to your child's doctor about additional vaccines that he may need.



U.S. Department of Health and Human Services
Centers for Disease Control and Prevention



American Academy of Pediatrics



DEDICATED TO THE HEALTH OF ALL CHILDREN™