

Global macroeconomic burden of epilepsy and the role for neurosurgery: a modelling study based upon the 2016 Global Burden of Disease data

T. J. O'Donohoe^{a,b,c}, A. Choudhury^b and E. Callander^b

^aDepartment of Neurosurgery, St Vincent's Hospital, Fitzroy, Vic.; ^bAustralian Institute of Tropical Health and Medicine, James Cook University, Townsville, QLD; and ^cNational Trauma Research Institute, Prahran, Vic., Australia

Keywords:

cost, economics, neurological conditions, neurosurgery, value of lost economic welfare

Received 28 April 2019
Accepted 3 September 2019

European Journal of Neurology 2020, **27**: 360–368

doi:10.1111/ene.14085

Background and purpose: Although the significant morbidity and mortality burden associated with epilepsy is well understood, associated economic consequences are yet to be estimated on a global scale. We sought to: (i) estimate the value of lost economic welfare attributable to epilepsy among countries included in the 2016 Global Burden of Disease study, (ii) evaluate differences in disease burden between countries of varied income classification and location, and (iii) understand the proportion of this burden that requires neurosurgical consultation and intervention.

Methods: Publicly available morbidity and mortality data were incorporated into a 'full-income' model to generate estimates of the cumulative value of lost economic welfare (VLW) related to epilepsy. Results from a survey of neurosurgeons were then used to estimate the VLW attributable to the proportion of disease requiring neurosurgical consultation and intervention.

Results: A total of 195 countries and territories were included in this analysis. We estimate that the cumulative VLW related to epilepsy was \$647.37 billion [2016 US dollars (USD), purchasing power parity (PPP)]. Economic welfare losses were equivalent to a mean of 1.45% ($\pm 1.00\%$) of gross domestic product. The value of economic losses attributable to the proportion of the burden necessitating neurosurgical consultation and intervention was \$258.95 billion (2016 USD, PPP) and \$155.37 billion (2016 USD, PPP) respectively.

Conclusions: Our results indicate that the economic consequences of epilepsy-related morbidity and mortality are substantial. When considered with evidence supporting the cost-effectiveness of various interventions for improved epilepsy diagnosis and management, our findings suggest that the implementation of simple and affordable measures may avert significant economic loss.

Introduction

Epilepsy is one of the most common chronic neurological conditions, directly affecting approximately 70 million people worldwide [1]. A recent estimate has suggested that this number is increasing by 4.6 million incident cases per year [2]. The vast majority of people with epilepsy reside in low-income countries (LICs)

and middle-income countries (MICs) [1,3], where the associated morbidity and mortality are significantly greater [1,4]. Although most patients do not receive appropriate treatment [5,6], between 70% and 85% will achieve remission with appropriately selected antiepileptic medications alone [7,8]. The remainder, however, continue to experience seizures and it has been suggested that these patients account for approximately 80% of the total direct healthcare costs attributable to epilepsy [9]. Surgical intervention has been demonstrated to be an efficacious and cost-effective intervention for a proportion of patients experiencing

Correspondence: T. J. O'Donohoe, St Vincent's Hospital, 41 Victoria Parade, Fitzroy, 3065 Vic., Australia (tel.: +61 (0) 3 9231 2211; fax: +61 (0) 3 9231 4871; e-mail: tomjmodonohoe@gmail.com).

medically refractory epilepsy [7,10–12]. Despite this, access to epilepsy surgery remains poor, particularly in LICs and MICs [13].

In addition to associated morbidity and mortality, people with epilepsy are at greater risk of poor educational, vocational, psychological and social outcomes [14,15]. Previous studies have adopted a number of perspectives from which to estimate the direct and indirect economic impact of these consequences [16]. However, to date, there has not been any published assessment of the global macroeconomic burden of epilepsy. When taken alongside cost-effectiveness data, such estimates could be utilized to inform decision-making regarding resource allocation and priority setting. With this in mind, we sought to: (i) estimate the total value of lost economic welfare (VLW) attributable to epilepsy at the global, regional and national level among the 195 countries and territories included in the Institute for Health Metrics and Evaluation (IHME) 2016 Global Burden of Disease (GBD) study [17], (ii) evaluate differences in disease burden between countries of varied income classification cohorts and locations, and (iii) understand the proportion of this burden that requires neurosurgical consultation and intervention. The counterfactual scenario in this analysis was the absence of disease.

Methods

This study has been reported according to the Guidelines for Accurate and Transparent Health Estimates Reporting [18]. A detailed description of the construction of our model can be found in Appendix S1 and a worked example can be found in Appendix S2.

Data sources

Burden of disease estimates and life expectancies were obtained from the IHME 2016 GBD study [17]. Gross domestic product (GDP) values for 2016 were extracted from the World Bank, measured in US dollars (USD) and adjusted for purchasing power parity (PPP) [19]. All countries were stratified into LICs, lower middle-income (LMIC), upper middle-income (UMIC) and high-income (HIC) countries according to the World Bank's classification system [20]. Sociodemographic index (SDI) values were obtained from the IHME.

Given that their estimates are derived from a comprehensive meta-analysis, we utilized the Value of a Statistical Life (VSL) published by the US Environmental Protection Agency for 2006, updated to the year of analysis (\$8 932 431).

Inclusion and exclusion criteria

Data from all countries included in the IHME 2016 GBD study were eligible for inclusion in this research. GDP data were not available for 11 of the included countries and territories (American Samoa, Andorra, Bermuda, Cuba, Djibouti, Eritrea, Greenland, Guam, North Korea, Northern Mariana Islands and Palestine), which precluded the calculation of a country-specific VSL. Given that these regions constituted such a small proportion of the overall population (0.66%), we assumed that these data were 'missing at random' and linearly projected existing estimates to encompass the entire target population. Disaggregated morbidity and mortality data were not available beyond 89 years of age and therefore the model was capped at this age.

Estimation of surgical burden

In the absence of quantitative, context-specific data to describe the surgical burden of epilepsy, the results of a survey were utilized to generate our estimates [21]. This survey involved 85 neurosurgeons from 57 countries who were asked to estimate the proportion of epilepsy that should, in ideal circumstances, receive neurosurgical consultation and intervention, respectively. Mean values supplied by the survey instrument suggested that 40% of epilepsy cases require neurosurgical consultation and 24% require neurosurgical intervention.

Sensitivity analyses

The sensitivity of our model to uncertainty intervals published by the IHME and the use of discounting were both evaluated with sensitivity analyses. Similarly, given the debate surrounding the relationship between VSL and income [22,23], the sensitivity of our estimates to variations in Income Elasticity (IE)-VSL was also investigated. Consistent with previous work by Alkire *et al.* [24] we utilized an IE-VSL of 0.55 for HICs and UMICs and 1.0 for LMICs and LICs as an upper bound, and an IE-VSL of 1.0 for HICs and UMICs and 1.5 for LMICs and LICs as a lower bound for these analyses. The estimate of epilepsy burden mandating neurosurgical intervention was also subjected to sensitivity analysis utilizing prevalence estimates of pharmaco-resistant disease of 22.5% and 37%, which were derived from large Scottish and French populations respectively [25,26].

Statistical analysis

Descriptive categorical data were summarized using absolute values and proportions, with numerical data

expressed with the mean (\pm SD). Comparisons of central tendency between normally distributed, independent groups were undertaken with a one-way ANOVA. Tukey and Games–Howell *post hoc* analyses were conducted when the assumption of homogeneity of variances was met, and violated, respectively.

Where data were normally distributed, without outlier values, the presence and direction of a linear relationship between continuous variables was determined using the Pearson product-moment correlation. The strength of the association was described according to Cohen, with coefficient values of 0.1–0.3 considered small in strength, 0.3–0.5 described as moderate and >0.5 described as strong [27].

Statistical analysis was conducted with Excel 2016 (Microsoft Corporation, Redmond, WA, USA) and Statistical Package for the Social Sciences Version 24 (IBM Corporation, Armonk, NY, USA). Graphs and figures were produced using Tableau (Tableau Software Inc., Seattle, WA, USA). A *P*-value of < 0.05 was considered to be statistically significant. All monetary estimates are expressed in terms of 2016 USD, adjusted for PPP.

Results

A total of 195 countries and territories were included in this analysis, with an estimated cumulative population of 7 434 043 990, which accounts for 99.86% of the global population in 2016. The mean GDP of included regions was \$19 212.11 (\pm \$20 599.59) and over 99% of countries and territories from each of the

World Bank income classifications were included in the model.

We estimate that the cumulative VLW related to epilepsy in 2016 was \$647.37 billion. The short-run impact of morbidity accounted for the majority (\$494.81 billion, 76.43%) of the overall burden of disease, whereas the long-term effects of mortality were responsible for \$152.55 billion (23.57%) of lost economic welfare. There was no difference in the proportion of lost welfare attributable to mortality between income [Welch's *F* (3.00, 87.49) = 1.72, *P* = 0.17] or geographic classification [Welch's *F* (6.00, 12.85) = 1.15, *P* = 0.39] groups. The age-specific burden of lost economic welfare is presented in Fig. 1.

Economic welfare losses were equivalent to a mean of 1.45% of GDP (\pm 1.00%). There was a statistically significant difference in VLW as a percentage of GDP between income classification groups [Welch's *F* (3.00, 82.44) = 40.56, *P* < 0.0005]. *Post hoc* analyses revealed significant differences between LICs and HICs (1.31; 95% confidence interval, 0.84–1.78; *P* < 0.0005), LMICs and HICs (1.19; 95% confidence interval, 0.77–1.61; *P* < 0.0005) and UMICs and HICs (0.98; 95% confidence interval, 0.60–1.35; *P* < 0.0005).

A similar difference was observed in the ratio of VLW:GDP between geographic groups [Welch's *F* (6.00, 25.81) = 14.37, *P* < 0.0005]. The mean ratio of VLW:GDP was highest in the Sub-Saharan Africa (2.20% \pm 1.08%) and South Asia (2.19% \pm 0.65%) groups, followed by the Middle East and North Africa (1.22% \pm 0.79%), East Asia and Pacific




Figure 1 Total value of lost economic welfare (VLW) [2016 US dollars (USD), purchasing power parity (PPP)] stratified by age group. [Colour figure can be viewed at wileyonlinelibrary.com]

(1.20% ± 1.00%), Europe and Central Asia (1.18% ± 0.91%), Latin America and the Caribbean (1.17% ± 0.51%) and North America (0.28% ± 0.25%). The results of *post hoc* analysis are presented in Table 1 and the global variation in VLW expressed as a proportion of GDP is illustrated in Fig. 2, with raw values also included in Appendix S3.

A Pearson's product-moment correlation revealed a moderate, inverse relationship between the ratio of VLW:GDP and SDI ($r = -0.48$, $P < 0.001$). An inverse relationship of smaller magnitude was also observed between the ratio of VLW attributed to YLLs and SDI ($r = -0.24$, $P = 0.001$).

Estimate of surgical burden

The estimated VLW attributable to the proportion of the burden of epilepsy necessitating neurosurgical consultation and intervention was \$258.95 billion and \$155.37 billion, respectively (Fig. 3a and b).

Sensitivity analysis

Utilizing the upper and lower IE-VSL values applied by Alkire *et al.* [24], the VLW estimate ranged between \$552.61 billion and \$862.69 billion. The model was more sensitive to variations in burden of disease estimates according to the uncertainty intervals published by the IHME, with a lower bound VLW estimate of \$342.69 billion and an upper bound of \$1090.26 billion. When based upon undiscounted years of life lost figures, the overall VLW estimate was \$752.57 billion. Utilization of the estimates of pharmacoresistance provided by Kwan and Brodie [25] and Picot *et al.* [26] as a surrogate for surgical burden yielded a VLW that ranged between \$145.66 billion and \$239.53 billion (Fig.4).

Discussion

We estimate that the cumulative VLW related to epilepsy among the 195 included countries and territories in 2016 was \$647.37 billion. Economic welfare losses were equivalent to a mean of 1.45% of GDP (±1.00%) and a significant difference was observed in the ratio of VLW:GDP between income classification groups [Welch's F (3.00, 82.44) = 40.56, $P < 0.0005$]. By providing the first global estimate of the macroeconomic burden of epilepsy, these findings extend those reported in a recent systematic review by Allers *et al.* [16] which summarized and critically appraised 22 studies describing the direct and indirect economic burden associated with epilepsy from the societal, health system and household perspective in a number of contexts.

Table 1 *Post hoc* analysis comparing the ratio of value of lost economic welfare: gross domestic product between geographic groups

Region	Mean difference	95% confidence interval	Significance
Sub-Saharan Africa			
East Asia & the Pacific	1.00	0.28 to 1.72	0.001*
Europe & Central Asia	1.01	0.40 to 1.62	<0.0005*
Latin America & the Caribbean	1.03	0.48 to 1.58	<0.0005*
Middle East & North Africa	0.98	0.26 to 1.69	0.002*
North America	1.91	1.10 to 2.71	0.0005*
South Asia	0.006	-0.94 to 0.95	>0.99
South Asia			
East Asia & the Pacific	0.99	0.02 to 1.96	0.04*
Europe & Central Asia	1.01	0.08 to 1.93	0.03*
Latin America & the Caribbean	1.02	0.12 to 1.93	0.03*
Middle East & North Africa	0.97	0.034 to 1.94	0.05
North America	1.91	0.89 to 2.92	0.001*
North America			
East Asia & the Pacific	-0.91	-1.73 to -0.09	0.03*
Europe & Central Asia	-0.90	-1.71 to -0.09	0.03*
Latin America & the Caribbean	-0.88	-1.78 to 0.02	0.05
Middle East & North Africa	-0.93	-1.76 to -0.11	0.03*
Middle East & North Africa			
East Asia & the Pacific	0.02	-0.73 to 0.78	>0.99
Europe & Central Asia	0.03	-0.63 to 0.70	>0.99
Latin America & the Caribbean	0.05	-0.55 to 0.66	>0.99
Latin America & the Caribbean			
East Asia & the Pacific	-0.03	-0.64 to 0.58	>0.99
Europe & Central Asia	-0.02	-0.49 to 0.45	>0.99
Europe & Central Asia			
East Asia & the Pacific	-0.01	-0.68 to 0.65	>0.99

The asterisk was used to reflect statistical significance.

In most circumstances, VLW estimates in HICs will exceed those in LICs by virtue of the more substantial VSLs found in wealthier economies. This can lead to the spurious conclusion that the macroeconomic burden of disease is greater in HICs than in their low- and middle-income counterparts. Therefore, although it is necessary to understand that our VLW estimates do not represent actual losses in GDP, their expression in this context provides an important sense of scale. The greater VLW:GDP ratio observed among LICs and MICs is a function of the greater burden of epilepsy found in regions with lower economic productivity, which is multifactorial in aetiology. Most epidemiological studies suggest that the prevalence of epilepsy is highest in resource-poor settings [2,17] and that focal-onset epilepsy is disproportionately




Figure 2 Value of lost economic welfare (VLW) expressed as a proportion of gross domestic product (GDP), stratified by country. [Colour figure can be viewed at wileyonlinelibrary.com]

represented when compared with HICs [28,29]. This observation supports the suggestion that the greater morbidity and mortality burden of epilepsy in LICs and LMICs reflects increased exposure to potentially epileptogenic insults, such as birth trauma, central nervous system infections, head injury and micronutrient deficiencies. The considerable incidence and prevalence of disease is compounded by a treatment gap of approximately 75% in many resource-poor countries [5,6]. Although the cause of this undertreatment varies according to prevailing sociocultural, geopolitical and economic conditions, inadequate access to skilled manpower, prohibitive treatment costs, poor availability of medications or diagnostic services, entrenched cultural beliefs regarding epilepsy causation and a preference toward traditional healers for treatment have all been identified as contributing factors [5,6].

Although mortality accounted for a minority [\$152.55 billion (23.57%)] of lost economic welfare, an inverse relationship, of modest magnitude, was observed between the ratio of VLW attributed to YLLs and SDI ($r = -0.24$, $P = 0.001$). This is consistent with evidence suggesting that the standardized mortality ratio in people with epilepsy in resource-poor settings such as Uganda (7.2) [30] and China (4.0) [31] significantly exceeds that in developed countries

(1.6–3.0) [32]. It is likely that this is attributable to the significant burden of untreated epilepsy discussed above and may also relate to an increased risk of fatal seizure complications such as drowning, burns and falls among those with poorer health literacy [33].

A number of studies have utilized similar methodology to estimate the VLW associated with various diseases [24,34–36]. In 2015, Alkire and colleagues calculated the present value of economic losses attributable to the cumulative surgical burden of injury, neoplasm, digestive disease and maternal and neonatal disorders to be \$14.5 trillion (2010 USD, PPP), equivalent to 17% of 2010 GDP [24]. More recently, Rudolfson *et al.* estimated that the VLW associated with neurosurgical disease (excluding epilepsy) in 2015 was \$3 trillion (2013 USD, PPP) [34]. Two smaller studies reported that the VLW associated with head and neck cancer in three South Asian countries, and the surgical proportion of cleft-palate repair in Sub-Saharan Africa was \$16.9 billion (2010 USD, PPP) [35] and 5.4–9.7 billion (2008 USD, PPP) [36], respectively. Although the differing sample populations, years of analysis and incorporated assumptions preclude direct comparison between our results, it is reassuring that they do not appear grossly disproportionate to those reported in this study.




Figure 3 (a) Total value of lost economic welfare (VLW) attributable to epilepsy necessitating neurosurgical consultation. (b) Total VLW attributable to epilepsy necessitating neurosurgical intervention. PPP, purchasing power parity; USD, US dollars. [Colour figure can be viewed at wileyonlinelibrary.com]

Surgical burden

Based upon the results of a survey involving neurosurgeons from all World Health Organization regions, we estimate that the total VLW attributable to the proportion of epilepsy necessitating neurosurgical consultation and intervention was \$258.95 billion and \$155.37 billion, respectively. Despite this significant burden, epilepsy surgery is globally underutilized [13], particularly in LICs and MICs [37,38]. The nature and causes of this underutilization have been comprehensively discussed elsewhere [13] and vary according to local socioeconomic, cultural and geopolitical circumstances. Broadly speaking, however, the increased penetration of epilepsy surgery centres relies upon the establishment of accessible medical infrastructure and expertise required to both evaluate patients for surgery and perform the necessary procedures. Clearly,

given that 75% of people with epilepsy do not receive appropriate medical treatment in some contexts [5], it is not justifiable to recommend universal scaling-up of surgical epilepsy services. Nevertheless, as the treatment gap continues to narrow, epilepsy surgery services should be considered as an important element of a comprehensive epilepsy management strategy, particularly for those patients with medically refractory epilepsy.

Strengths and limitations

It is important that our estimates be interpreted in the knowledge of their limitations. Although we believe that it provides the fullest measure of total economic welfare losses, there are empirical concerns about the ‘full-income’ model, with many proposing that the derivation of VSL from willingness-to-pay estimates is




Figure 4 Sensitivity analysis demonstrating estimates of value of lost economic welfare (VLW) based upon upper and lower Income Elasticity of the Value of a Statistical Life (IE-VSL) estimates, upper and lower bound Institute for Health Metrics and Evaluation (IHME) estimates and undiscounted years of life lost (YLL). PPP, purchasing power parity; USD, US dollars. [Colour figure can be viewed at wileyonlinelibrary.com]

likely to be oversimplistic [39]. Similarly, given that it places the largest value on each disability-adjusted life-year incurred, this methodology consistently generates larger estimates of economic loss than similar cost-of-illness methodologies. We must also acknowledge the significant role played by various assumptions in the generation of estimates in the IHME GBD study and in our model. The impact of the most salient of these assumptions have been interrogated with sensitivity analysis and this is supplemented with a detailed explanation of the rationale for each of the assumptions in order to permit appraisal of the validity of our estimates. Similarly, although there will probably be some debate regarding the estimates of disease burden mandating neurosurgical consultation and intervention, in the absence of more compelling information from a range of settings, we thought it reasonable to derive these proportions from a geographically and socioeconomically representative sample of neurosurgeons [21], which is consistent with previous similar studies [24,34–36]. We also acknowledge that the application of a uniform proportion of surgical burden to all countries is likely to be simplistic. Indeed, given that focal epilepsy appears to be more prevalent in LICs and LMICs, it is possible that epilepsy surgery has a greater role in these settings than it does in more developed countries, although this proposal remains conjectural and any attempt to account for it in this analysis would be speculative. GDP data were missing for a small number of countries, which prohibited the derivation of a country-specific VSL and, thus, estimation of VLW. Although some may protest their inclusion in the headline

figures, this had minimal impact on our overall estimate (0.62%). Finally, it is vital to emphasize the limitations of conclusions that may be reached from economic impact studies. Indeed, although our estimate of the macroeconomic burden of epilepsy may be valuable, these data must be validated with context-specific estimates and considered alongside cost-effectiveness data before making allocative decisions.

In spite of these limitations, which are commonplace in much of the health economic literature, this study is notable because it utilizes a commonly accepted economic methodology, which incorporates both market and non-market losses, to provide the first global macroeconomic estimate of the burden of lost economic welfare related to epilepsy and the proportion that may be amenable to neurosurgical consultation or intervention.

Conclusion

The significant morbidity and mortality burden of epilepsy has been clearly established in the IHME 2016 GBD study [17]. Consistent with these findings, our results indicate that the cumulative value of associated economic welfare losses is substantial. After adjustment for GDP, it is clear that this economic impact disproportionately affects resource-poor countries. When considered alongside evidence supporting the cost-effectiveness of various interventions for improved epilepsy diagnosis and management in countries of all income classifications, our findings suggest that the implementation of simple and affordable interventions may avert significant economic loss. We

also suggest that the value of economic losses attributable to cases of epilepsy requiring neurosurgical consultation or intervention is considerable. Although the prioritization of epilepsy surgery services cannot be advocated in all settings, it is clear that cost-effective surgical interventions can be delivered in resource-limited countries and should have an increasing role in epilepsy management strategies worldwide. We hope that these estimates act as a stimulus for further research and collaboration between neurologists, neurosurgeons and policy-makers to better understand and respond to the diverse challenges facing cost-effective epilepsy care across the world.

Supporting Information

Additional Supporting Information may be found in the online version of this article:

Appendix S1. Conceptual Overview of Approach & Model Construction.

Appendix S2. Worked Example – Afghanistan (<12 months).

Appendix S3. Raw headline figures for VLW/GDP (%).

References

- Ngugi AK, Bottomley C, Kleinschmidt I, Sander JW, Newton CR. Estimation of the burden of active and lifetime epilepsy: a meta-analytic approach. *Epilepsia* 2010; **51**: 883–890.
- Vaughan KA, Lopez Ramos C, Buch VP, *et al.* An estimation of global volume of surgically treatable epilepsy based on a systematic review and meta-analysis of epilepsy. *J Neurosurg* 2018; **130**: 1–15.
- Winkler AS, Schaffert M, Schmutzhard E. Epilepsy in resource poor countries – suggestion of an adjusted classification. *Epilepsia* 2007; **48**: 1029–1030.
- Diop AG, Hesdorffer DC, Logroscino G, Hauser WA. Epilepsy and mortality in Africa: a review of the literature. *Epilepsia* 2005; **46**(Suppl. 11): 33–35.
- Meyer AC, Dua T, Ma J, Saxena S, Birbeck G. Global disparities in the epilepsy treatment gap: a systematic review. *Bull World Health Organ* 2010; **88**: 260–266.
- Mbuba CK, Ngugi AK, Newton CR, Carter JA. The epilepsy treatment gap in developing countries: a systematic review of the magnitude, causes, and intervention strategies. *Epilepsia* 2008; **49**: 1491–1503.
- Picot MC, Jaussent A, Neveu D, *et al.* Cost-effectiveness analysis of epilepsy surgery in a controlled cohort of adult patients with intractable partial epilepsy: A 5-year follow-up study. *Epilepsia* 2016; **57**: 1669–1679.
- Brodie MJ, Barry SJ, Bamagous GA, Norrie JD, Kwan P. Patterns of treatment response in newly diagnosed epilepsy. *Neurology* 2012; **78**: 1548–1554.
- Begley CE, Famulari M, Annegers JF, *et al.* The cost of epilepsy in the United States: an estimate from population-based clinical and survey data. *Epilepsia* 2000; **41**: 342–351.
- Engel J Jr, McDermott MP, Wiebe S, *et al.* Early surgical therapy for drug-resistant temporal lobe epilepsy: a randomized trial. *JAMA* 2012; **307**: 922–930.
- Wiebe S, Blume WT, Girvin JP, Eliasziw M. A randomized, controlled trial of surgery for temporal-lobe epilepsy. *N Engl J Med*. 2001; **345**: 311–318.
- West S, Nolan SJ, Cotton J, *et al.* Surgery for epilepsy. *Cochrane Database Syst Rev* 2015; Cd010541. <https://doi.org/10.1002/14651858.CD010541.pub2> [Epub ahead of print].
- Jette N, Sander JW, Keezer MR. Surgical treatment for epilepsy: the potential gap between evidence and practice. *Lancet Neurol* 2016; **15**: 982–994.
- Salpekar JA, Mula M. Common psychiatric comorbidities in epilepsy: How big of a problem is it? *Epilepsy Behav* 2018. in press.
- Laxer KD, Trinka E, Hirsch LJ, *et al.* The consequences of refractory epilepsy and its treatment. *Epilepsy Behav* 2014; **37**: 59–70.
- Allers K, Essue BM, Hackett ML, *et al.* The economic impact of epilepsy: a systematic review. *BMC Neurol* 2015; **15**: 245.
- GBD 2016 DALYs and HALE Collaborators. Global, regional, and national disability-adjusted life-years (DALYs) for 333 diseases and injuries and healthy life expectancy (HALE) for 195 countries and territories, 1990–2016: a systematic analysis for the Global Burden of Disease Study 2016. *Lancet* 2017; **390**: 1260–1344.
- Stevens GA, Alkema L, Black RE, *et al.* Guidelines for Accurate and Transparent Health Estimates Reporting: the GATHER statement. *Lancet* 2016; **388**: e19–e23.
- The World Bank Group. <https://data.worldbank.org/indicator/ny.gdp.mktp>. pp. cd (accessed 15/08/2017).
- Fantom N, Serajuddin U. The World Bank's Classification of Countries by Income. Policy Research Working Paper; No. 7528. Washington, DC: The World Bank, 2016.
- Dewan MC, Rattani A, Baticulon RE, *et al.* Operative and consultative proportions of neurosurgical disease worldwide: estimation from the surgeon perspective. *J Neurosurg* 2019; **130**: 1098–1106.
- Viscusi WK, Aldy JE. The value of a statistical life: a critical review of market estimates throughout the world. *J Risk Uncertain* 2003; **27**: 5–76.
- Hammit JK, Robinson LA. The income elasticity of the value per statistical life: transferring estimates between high and low income populations. *J Benefit Cost Anal* 2011; **2**: 1–29.
- Alkire BC, Shrimpe MG, Dare AJ, Vincent JR, Meara JG. Global economic consequences of selected surgical diseases: a modelling study. *Lancet Glob Health* 2015; **3** (Suppl. 2): S21–27.
- Kwan P, Brodie MJ. Early identification of refractory epilepsy. *N Engl J Med* 2000; **342**: 314–319.
- Picot MC, Baldy-Moulinier M, Daures JP, Dujols P, Crespel A. The prevalence of epilepsy and pharmacoresistant epilepsy in adults: a population-based study in a Western European country. *Epilepsia* 2008; **49**: 1230–1238.
- Cohen J. *Statistical Power Analysis for the Behavioural Sciences*, 2nd edn. New Jersey, Lawrence Erlbaum, 1988.
- Mac TL, Tran DS, Quet F, Odermatt P, Preux PM, Tan CT. Epidemiology, aetiology, and clinical

- management of epilepsy in Asia: a systematic review. *Lancet Neurol* 2007; **6**: 533–543.
29. Preux PM, Druet-Cabanac M. Epidemiology and aetiology of epilepsy in sub-Saharan Africa. *Lancet Neurol* 2005; **4**: 21–31.
 30. Kaiser C, Asaba G, Kasoro S, Rubaale T, Kabagambe G, Mbabazi M. Mortality from epilepsy in an onchocerciasis-endemic area in West Uganda. *Trans R Soc Trop Med Hyg* 2007; **101**: 48–55.
 31. Mu J, Liu L, Zhang Q, *et al.* Causes of death among people with convulsive epilepsy in rural West China: a prospective study. *Neurology* 2011; **77**: 132–137.
 32. Forsgren L, Hauser WA, Olafsson E, Sander JW, Sillanpaa M, Tomson T. Mortality of epilepsy in developed countries: a review. *Epilepsia* 2005; **46**(Suppl. 11): 18–27.
 33. Beghi E. Injuries in patients with epilepsy. *Lancet Neurol* 2005; **4**: 71–72.
 34. Rudolfson N, Dewan MC, Park KB, Shrimel MG, Meara JG, Alkire BC. The economic consequences of neurosurgical disease in low- and middle-income countries. *J Neurosurg* 2019; **130**: 1149–1156.
 35. Alkire BC, Bergmark RW, Chambers K, *et al.* Head and neck cancer in South Asia: Macroeconomic consequences and the role of the head and neck surgeon. *Head Neck* 2016; **38**: 1242–1247.
 36. Corlew DS, Alkire BC, Poenaru D, Meara JG, Shrimel MG. Economic valuation of the impact of a large surgical charity using the value of lost welfare approach. *BMJ Global Health* 2016; **1**(4): e000059.
 37. Dua T, de Boer HM, Prilipko LL, Saxena S. Epilepsy Care in the World: results of an ILAE/IBE/WHO Global Campaign Against Epilepsy survey. *Epilepsia* 2006; **47**: 1225–1231.
 38. Qiu J. Epilepsy surgery: challenges for developing countries. *Lancet Neurol* 2009; **8**: 420–421.
 39. World Health Organisation. WHO Guide to Identifying the Economic Consequences of Disease and Injury. Department of Health Systems Financing. Geneva, Switzerland: World Health Organisation. 2009.