

Two C-Diff Best Practice Advisories (BPAs) Will Be Reactivated

Effective Tuesday, August 2, two C-Diff BPAs will be reactivated in Production:

- Fire BPA alert when C-Diff order is placed on a patient with laxatives, tube feeds, or oral contrast in the last 48-hours.

BestPractice Advisory - Nouwick, Fleur

Important (1)

⚠ C Diff testing is contraindicated on this patient due to the following:

- ✓ Laxatives administered in the last 48 Hours

For clinically stable patients (e.g. patients without fever, abdominal pain/distention, or leukocytosis), please wait 48 hours after last administration prior to assessing for ongoing diarrhea. If patient is taking a daily laxative please consider whether patient's stool frequency or consistency is worse than their baseline.

Click '**Accept**' to remove the order from 'Order Entry'.

Remove the following orders? _____

Remove	Keep	
<input checked="" type="checkbox"/>	<input type="checkbox"/>	Clostridium difficile PCR Once, today at 1619, For 1 occurrence Stool, Per Rectum

Acknowledge Reason _____

Concern for severe disease; diagnosis ca...	Concern for ileus due to C. difficile	
Pt has been on laxatives & diarrhea was ...	Pt has been on tube feeds & diarrhea was...	Other (Comment)

☒ **Accept**

- Fire BPA alert when C-Diff order is placed on a patient with an existing order, a previous positive in the last 14-days, or a previous negative in the last 7-days.

BestPractice Advisory - Wiggum, Clancy

Important (1)

ⓘ C Diff testing is contraindicated on this patient due to the following:

✔ Positive C diff result in last 14 days

Click **'Accept'** to remove the order from 'Order Entry'.

Remove the following orders?

Remove

Keep

Clostridium difficile PCR

Once, today at 1012, For 1 occurrence Stool, Per Rectum

Acknowledge Reason

Reason for Ordering

✔ Accept

Updated 8/1/2023.

2