


Orders: Admission Medication Reconciliation


Medication Reconciliation (Med Rec) is the process of comparing a patient’s current orders or home medication list with new orders you are writing. To maintain a complete and accurate list of the medications, reconcile medications at the following times: Admission, Transfer and Discharge.

Sequence of steps in hospital admission med reconciliation:




How it Works


1. Select the Admission Navigator and then select Admission orders from the side menu
 - NOTE: "Admission" may be truncated to "Admis..." or "Adm..." depending on the width of your screen




2. Select the tab "Review Home Medications"




3. If there is an alert that "Medications from outside sources need attention" click on "Go Reconcile" first




- Because Epic is receiving data from multiple heritage systems, from CareEverywhere and from Health Information Exchanges, there are likely to be many outside medications to be reviewed.
- Reasons to do this now are:
 1. The patient's home medication list in Epic may not have the most current information.
 2. The outside sources may include newer medications or dosage adjustments that were created outside of Epic.
 3. The outside sources are likely to include duplicates of medications on the patient's Epic medication list.
 4. Good citizenship: resolving any discrepancies at this stage will spare future users from uncertainty about what the patient is actually taking.




- Medications from outside sources will be marked with a red "e".
- In this case, "Aspirin" is a duplicate of a medication already on the patient's medication list, so click on the trashcan icon to delete it.



- For medications that should be added to the current home medication list, click on the "+" icon, instead.
 - When finished, click on the "Accept" button to save any changes.
4. Add any medications the patient is taking that are not listed here by selecting the "Add Prior to Admission Med" field.





- For each medication, click on the appropriate button indicating the timing of the most recent dose.
- If the patient is taking the medication, click on the "Taking" box. If the patient is NOT taking the medication, leave the box blank.



NOTE: Reconciliation of home medications may be carried out only by a licensed independent provider: Physician (including residents and fellows), APN, or PA.


5. Select the "Reconcile Home Medications" tab.



6. Until this process is complete, there will be a warning in the side bar.

7. For each medication, indicate the appropriate action:

- **Order:** Will be given in hospital
- **Don't Order:** Will not be given in hospital
- **Replace:** Order an alternate med
- **Remove:** Discontinue, including from home medication list
- **Order and Hold:** Will be ordered, but will not be given until the hold is manually released.



8. For some medications, you will have to specify an alternative because the home medication is not available on the hospital formulary

- If that is the case, you will be given a list of the available alternatives to the home medication

Alternative Required

You are reordering:
omeprazole (PriLOSEC) 20 mg DR capsule: Take by mouth 1 (one) time each day., Starting Tue 1/18/2022, Historical Med

Details

Pantoprazole is the P&T approved formulary substitution for Proton Pump Inhibitors (PPIs).
Choose one of the substitutions below.

Original Medication	Original Dosage	Substitution Medication	Substitution Dosage
ORAL (PO)			
dexlansoprazole (DEXILANT)	30mg	pantoprazole (PROTONIX)	20mg Daily
dexlansoprazole (DEXILANT)	60mg	pantoprazole (PROTONIX)	40mg Daily
esomeprazole (NEXIUM)	20mg Daily	pantoprazole (PROTONIX)	40mg Daily
esomeprazole (NEXIUM)	40mg Daily	pantoprazole (PROTONIX)	40mg BID

Alternatives

Alternative	Details	Cost
<input type="radio"/> pantoprazole (ProtoNix) EC tablet	20 mg, oral, Daily before breakfast	\$
<input type="radio"/> pantoprazole (PROTONIX) EC tablet	40 mg, oral, Daily before breakfast	\$
<input type="radio"/> pantoprazole (ProtoNix) EC tablet	40 mg, 2 times daily before meals	\$
<input type="radio"/> pantoprazole (PROTONIX) packet	40 mg, nasogastric tube, Daily before breakfast	
<input type="radio"/> pantoprazole (ProtoNix) packet	40 mg, nasogastric tube, 2 times daily before meals	
<input type="radio"/> pantoprazole (PROTONIX) IV	40 mg, intravenous, Daily	\$\$

Accept Alternative
 Cancel

Version #	Purpose of Change	Author	Date of Change
1.0	Initial Creation	M. Bauerlein	5/13/22
1.1	Updated	M. Bauerlein	5/16/22