



Grazing & Hay Restrictions for Pasture & Rangeland Herbicides



Herbicides	Product	Lactating Dairy Animals		Beef and Non-Lactating Dairy Animals		
		Before Grazing	Before Hay Harvest	Before Grazing	Before Hay Harvest	Removal Before Slaughter
Amber	0.28 – 0.56 oz	0	0	0	0	0
Chaparral	1.0 – 3.3 oz	0	0	0	0	0
Cimarron Plus	0.125-1.25 oz	0	0	0	0	0
Crossbow L	1 to 6 qt	1 year	14 days	0	0	3 days
Curtail	2 – 4 qt	14 days	7 days	0	30 days	7 days(G)
Dicamba (Rifle)	Up to 1 pt	7 days	37 days	0	0	30 days
	Up to 2 pt	21 days	51 days	0	0	30 days
	Up to 4 pt	40 days	70 days	0	0	30 days
Distinct	4.0 to 8.0 oz	0	0	0	0	0
DuraCor (H)	12 to 20 fl oz	0	0	0	0	0
Escort XP	0.33-1.66 oz	0	0	0	0	0
	1.67-3.33 oz	3 days	3 days	3 days	3 days	3 days
ForeFront HL	1.2 – 2.1 pts	0	7 days	0	7 days	0
Garlon 3A	0.5 – 1.5 gal/100 gal	1 year	1 year	14 days	14 days	3 days
Gramoxone SL(B)	1.5-2.0 pt	1 month	1 month	1 month	1 month	0
Grazonnext HL	Up to 34 fl oz	0	7 days	0	7 days	0
Grazon P+D (C)	3.0 – 4.0 pt	7 days	30 days	0	30 days	3 days
Milestone	3.0 – 7.0 oz	0	0	0	0	0
Overdrive	4.0 – 8.0 oz	0	0	0	0	3 days
PastureGard HL	1.0 to 1.5 pts	1 year	1 year	0	14 days	3 days
Plateau	4.0-12.0 oz	0	7 days	0	7 days	0
Rave	2.0-5.0 oz	7 days	-	0	-	30 days
Redeem R&P	1.5 – 4.0 pt	14 days	365 days	0	7 days	3 days
Remedy	1.0 – 2.0 qt	14 days	1 year	0	14 days	3 days
Rifle-D	2.0 – 4.0 pts	7 days	37 days	0	0	30 days
Roundup (see individual labels, as they may differ)						
Spot or wiper(A)	Any labeled rate	14 days	14 days	14 days	14 days	0
Broadcast	Any labeled rate	8 weeks	8 weeks	8 weeks	8 weeks	0
Saber	½ to 4 pts	7 days	30 days	0	30 days	3 days
Salvo	12.8 – 25.6 oz (up to 51.2 oz for brush)	7 days	30 days	0	30 days	3 days
Sharpen	1-2 oz	0	0	0	0	0
Spike 20P	½ oz per 45 sq. ft.	0 days (G)	1 year (F)	0	1 year (F)	0 days (F)
Surmount	1.5 – 4.0 pt	14 days	7 days	0	7 days	3 days
Stinger 3E	0.66 – 1.31 pt	0	0	0	0	0
Telar XP	Up to 1.33 oz	0	0	0	0	0
Tordon 22K (C)	0.5 – 2.0 pt	14 days	14 days	0	14 days	3 days
Transline	0.25 – 1.3 pt	0	0	0	0	0
2,4-D/MCPA(D)	0.5 pt – 1.0 qt	7 – 14 days	30 days	0 – 7 days	0 – 30 days	0

(A)Do not treat more than 1/10th of any given acre at one time with spot or wiper applications. Remove livestock before application.

(B)Restrictions based on the degree of new seedling establishment before grazing. Suggested at least 6" of grass or legume seedling growth, which is approximately 1 month. Late fall seeding may require 3- 5 months before the suggested 6" height is reached.

(C)Move livestock to untreated grass pasture for 7 days before transferring livestock to broadleaf crop or pasture areas. Removal for slaughter statement only applies to animals grazing treated area forage for 2 weeks immediately after application. Do not exceed 1 qt/A broadcast. If >1qt/A; only spot treatments are allowed; total acres cannot exceed 25% of area in any particular watershed.

(D)Be sure to check individual product labels for restrictions and use rates due to the large number of formulations available.

(E)One year if more than 1.5 gal/A rate used.

(F)If no more than 20 lb./A used.

(G)Withdrawal not needed if 2 weeks or more time elapsed since application.

(H) Do not use grasses treated in preceding 18 months for export or off the farm/ranch. Consult label/supplemental labels.

Always Read and Follow Label Directions & Precautions. This is just a guide!

This info was compiled using the Individual labels, Illinois Ag Pest Handbook, and Herbicide Use in Nebraska Guide. Errors or omissions may occur. Some labels list a difference between green or dry hay. This list is dry, not a green chop. Look to label for individual needs.