

Garden Tips for April

Fruit and Nut

- Don't spray insecticides during fruit tree bloom or pollination may be affected. Disease sprays can continue according to schedule and label directions. ([EPP-7319](#))
- Control cedar-apple rust. When the orange jelly galls are visible on juniper (cedar), following a rain, begin treating apple and crabapple trees with a fungicide. ([EPP-7319](#), [HLA-7611](#))
- Fire blight bacterial disease can be controlled at this time. Plant disease-resistant varieties to avoid diseases.
- Continue spray schedules for disease prone fruit and pine trees.

Tree and Shrub

- Proper watering of newly planted trees and shrubs often means the difference between success and replacement.
- Remove any winter-damaged branches or plants that have not begun to grow. Prune spring flowering plants as soon as they are finished blooming. ([HLA-6404](#), [HLA-6409](#))
- Control of powdery mildew disease can be done with early detection and regular treatment. Many new plant cultivars are resistant. ([EPP-7617](#))
- Leaf spot diseases can cause premature death of foliage and reduce plant vigor.

Flowers

- Most bedding plants, summer flowering bulbs, and annual flower seeds can be planted after danger of frost. This happens around mid-April in most of Oklahoma. Hold off mulching these crops until spring rains subside and soil temperatures warm up. Warm-season annuals should not be planted until soil temperatures are in the low 60s.
- Harden off transplants outside in partial protection from sun and wind prior to planting.
- Let spring flowering bulb foliage remain as long as possible before removing it.

Vegetables

- Wait a little longer for it to warm up before planting cucurbit crops and okra.
- Plant vegetable crops in successive plantings to ensure a steady supply of produce rather than harvesting all at once.
- Cover cucurbit crops with a floating row cover to keep out insect pests. Remove during bloom time.
- Watch for cutworm damage and add flea beetle scouting to your list of activities in the vegetable garden.

Garden Planting Guide for Warm-Season Vegetables

<u>Vegetable</u>	<u>Time to Plant*</u>	<u>Days to Harvest</u>	<u>Method of Planting</u>
Bean, Lima	April 15-30	90-120	Seed
Beans, Green or Wax	April 10-30	50-60	Seed
Beans, Pole	April 10-30	60-90	Seed
Cantaloupe	May 1-20	80-100	Seed or Plants
Cucumber	April 10-30 or later	50-70	Seed or Plants
Eggplant	April 10-30	80-90	Plants
Okra	April 10-30 or later	60-70	Seed
Pepper	April 10-30 or later	90-110	Plants
Pumpkin	April 10-30	90-120	Seed
Southern Pea	May 1-June 10	85-100	Seed
Squash, Summer	April 10-30 or later	40-60	Seed or Plants
Squash, Winter	May 15-June 15	110-125	Seed or Plants
Sweet Corn	Mar. 25-April 30	80-100	Seed
Sweet Potato	May 1-June 10	100-120	Plants
Tomato	April 10-30	70-90	Plants
Watermelon	May 1-20	90-120	Seed

**These dates indicate planting times from southeast to northwest Oklahoma. Specific climate and weather may influence planting dates. For Cool-Season Vegetables, the soil temperature at the depth where the seeds are planted should be at least 40°F.*

Landscape – General

- Hummingbirds arrive in Oklahoma in early April. Get your bird feeders ready using 1 part sugar to 4 parts water. Do not use red food coloring.
- Keep the bird feeder filled during the summer and help control insects at the same time.
- Lace bugs, aphids, spider mites, bagworms, etc. can start popping up in the landscape and garden later this month. Keep a close eye on all plants and use mechanical, cultural, and biological control options first.
- Be alert for both insect pests and predators. Some pests can be hand picked without using a pesticide. Do not spray if predators such as lady beetles are present. Spray only when there are too few predators to be effective.

Lawn

- Warm-season grass lawns can be established beginning late April from sprigs, plugs, or sod. ([HLA-6419](#))

- Fertilizer programs can begin for warm-season grasses in April. The following recommendations are to achieve optimum performance and appearance of commonly grown species in Oklahoma.

- Zoysiagrass: 3 lbs N/1,000 sq. ft./year
- Bahiagrass: 3 lbs N/1,000 sq. ft./year
- Buffalograss: 2 - 3 lbs N/1,000 sq. ft./year
- Buffalograss/grama mixes: 3 lbs N/1,000 sq. ft./year
- Bermudagrass: 4-6 lbs N/1,000 sq. ft./year
- Centipedegrass: 2 lbs N/1,000 sq. ft./year
- St. Augustinegrass: 3-6 lbs N/1,000 sq. ft./year

When using quick release forms of fertilizer, use 1 pound of actual nitrogen per 1,000 sq. ft. per application; water in nitrate fertilizers. ([HLA-6420](#))

- Mowing of warm-season lawns can begin now ([HLA-6420](#)). Cutting height for bermudagrass and zoysiagrass should be 1 to 1½ inches high, and buffalograss 1 ½ to 3 inches high.
- Damage from Spring Dead Spot Disease (SDS) becomes visible in bermudagrass ([EPP-7665](#)). Perform practices that promote grass recovery. Do not spray fungicides at this time for SDS control.
- Grub damage can be visible in lawns at this time. Check for the presence of grubs before ever applying any insecticide treatments. Apply appropriate soil insecticide if white grubs are a problem ([EPP-7306](#)). Water product into soil.