

PENNSYLVANIA DEPARTMENT OF HEALTH
 2026 – PAHAN –824-05-15 -ADV
Anticipated Seasonal Increase in *Legionella* Infections



Media Disclaimer: *The Pennsylvania Health Alert Network (PA-HAN) provides timely information on emerging public health issues to state and local public health agencies, hospitals, emergency management officials, and other health care providers. These messages are intended to inform and support clinical work and should not be treated as news releases. If members of the media have questions about a PA-HAN, contact the DOH Office of Communications at ra-dhpressoffice@pa.gov.*

DATE:	5/15/2026
TO:	Health Alert Network
FROM:	Debra L. Bogen, MD, FAAP, Secretary of Health
SUBJECT:	Anticipated Seasonal Increase in <i>Legionella</i> Infections
DISTRIBUTION:	Statewide
LOCATION:	Statewide
STREET ADDRESS:	n/a
COUNTY:	n/a
MUNICIPALITY:	n/a
ZIP CODE:	n/a

This transmission is a “Health Advisory”: provides important information for a specific incident or situation; may not require immediate action.

HOSPITALS: PLEASE SHARE WITH ALL MEDICAL, PEDIATRIC, NURSING AND LABORATORY STAFF IN YOUR HOSPITAL; **EMS COUNCILS:** PLEASE DISTRIBUTE AS APPROPRIATE; **FQHCs:** PLEASE DISTRIBUTE AS APPROPRIATE; **LOCAL HEALTH JURISDICTIONS:** PLEASE DISTRIBUTE AS APPROPRIATE; **PROFESSIONAL ORGANIZATIONS:** PLEASE DISTRIBUTE TO YOUR MEMBERSHIP; **LONG-TERM CARE FACILITIES:** PLEASE SHARE WITH ALL MEDICAL, INFECTION CONTROL AND NURSING STAFF IN YOUR FACILITY.

Summary

- In Pennsylvania, *Legionella* infections increase each year during spring, summer, and fall months (May-November).
- The preferred diagnostic tests for a *Legionella* infection are culture of lower respiratory secretions (e.g., sputum, bronchoalveolar lavage) on selective media AND the *Legionella* urinary antigen test. Submit *Legionella* isolates to the PADOH Bureau of Laboratories for characterization.
- Health care providers should maintain a high index of suspicion for Legionnaire’s disease in adult patients with pneumonia.
- Report all *Legionella* infections to the health department via PA-NEDSS or by calling PADOH (877-PA-HEALTH) or the local health department.

Background

Since 2000, the number of reported *Legionella* infections have been increasing in both Pennsylvania and the United States overall. *Legionella* infections follow a seasonal pattern. The majority of cases occur between May and November each year. In May and June of 2025, Pennsylvania *Legionella* infections were the highest observed in the last five years. *Legionella* bacteria grow best in warm, humid, and wet weather. In addition to seasonality and temperature, stagnant water due to building closures or dead legs in pipe networks may allow for *Legionella* growth. More information on environmental factors that

contribute to *Legionella* growth can be found on the CDC's website: <https://www.cdc.gov/control-legionella/php/public-health-strategy/index.html>

Many symptoms of *Legionella* infections are similar to other respiratory infections. Due to these similarities, clinicians may test certain patients presenting with respiratory symptoms for other respiratory pathogens and not for *Legionella* infection. Delayed diagnosis may slow appropriate treatment. For this reason, it is recommended that all patients hospitalized with pneumonia of unknown cause be tested for *Legionella* infection.

***Legionella* infection (legionellosis) can manifest as Legionnaires' disease, Pontiac fever, or extrapulmonary legionellosis.**

- **Legionnaires' disease** is a severe illness with pneumonia. Symptoms include cough, shortness of breath, fever, muscle aches, and headaches. Some patients also experience diarrhea, nausea, and confusion. Most patients are hospitalized, and treatment is required. The case-fatality rate is about 10% for community-acquired Legionnaires' disease and about 25% for healthcare-acquired disease.
- **Pontiac fever** is a milder illness, frequently characterized by fever and muscle aches. Patients with Pontiac fever do not develop pneumonia, do not require treatment, and typically recover within a week.
- **Extrapulmonary legionellosis** is a *Legionella* infection at a site outside the lungs (e.g., endocarditis, wound infection, joint infection, or graft infection). Symptoms and treatment vary according to site of infection.

Risk factors for *Legionella* infection include:

- Age ≥50 years
- Current or former smoker
- Underlying conditions such as chronic lung disease, cancer, diabetes, renal disease, or immunocompromising conditions
- Recent travel with an overnight stay outside of the home
- Recent care at a health care facility
- Exposure to hot tubs

A *Legionella* infection occurs when a person inhales aerosolized water containing *Legionella* bacteria. Potential sources include hot tubs, showers, faucets, decorative fountains, and cooling towers. Patients may also be infected through aspiration of contaminated drinking water. In general, people do not spread Legionnaires' disease and Pontiac fever to other people. Most cases are sporadic, but outbreaks can occur.

The incubation period for Legionnaires' disease is generally 5 to 6 days (range 2-14 days). For Pontiac fever, the incubation period is 1 to 3 days but can be as short as 4 hours.

Testing

The preferred diagnostic tests for *Legionella* infection are the *Legionella* urinary antigen test AND culture of lower respiratory secretions. Serologic assays can be nonspecific and are not recommended in most situations. Testing is warranted for the following indications:

- Patients who have failed outpatient antibiotic therapy for community-acquired pneumonia
- Patients with severe pneumonia, particularly those requiring intensive care
- Patients with pneumonia who are immunocompromised
- Patients who traveled away from their home in the 14 days prior to the onset of illness
- Patients with pneumonia who spent time in a location with a *Legionella* outbreak or a location with a known *Legionella* infection in the last 12 months.

- Patients at risk for *Legionella* infection with healthcare-associated pneumonia (pneumonia with onset ≥48 hours after admission)
- Patients with pneumonia who spent time in a location with a known positive *Legionella* water sample

Best practice is to obtain both the urinary antigen test and sputum culture concurrently. Sputum should ideally be obtained prior to antibiotic administration, but antibiotic treatment should not be delayed to facilitate this process.

- ***Legionella* urinary antigen test**

The most commonly used laboratory test for diagnosis of a *Legionella* infection is the urinary antigen test, which detects a molecule of the *Legionella* bacterium in urine. The test can remain positive for a few weeks after infection, even with antibiotic treatment. The urinary antigen test only detects the most common cause of *Legionella* infection, *L. pneumophila* serogroup 1. However, other species and serogroups of *Legionella* are pathogenic, so a patient with a negative urinary antigen result could have legionellosis caused by other *Legionella* species and serogroups.

- **Culture of lower respiratory secretions (e.g., sputum, bronchoalveolar lavage) on selective media**

Culture can detect *Legionella* species and serogroups that the urinary antigen test does not, and it allows for the comparison of clinical and environmental isolates in the event of an outbreak. When specimens are submitted for culture, laboratories must be informed that *Legionella* is suspected because it requires the use of specialized media (Buffered Charcoal Yeast Extract [BCYE] agar).

***Legionella* isolates should be forwarded to PADOH Bureau of Laboratories for further characterization.**

- Ship isolates at room temperature. Slanted media is preferred. If plates are sent, please seal with parafilm. Package as Category B sample and ship (Monday through Thursday only) to:

PA Department of Health, Bureau of Laboratories
 110 Pickering Way
 Exton, PA 19341
 610-280-3464

Treatment

For patients with Legionnaires' disease, follow the Infectious Diseases Society of America (IDSA)/American Thoracic Society (ATS) guidelines for the treatment of [community-acquired](#) and [hospital-acquired](#) pneumonia. *Legionella*-directed antibiotics include macrolides and respiratory fluoroquinolones. While it is preferred that diagnostic testing specimens are obtained before antibiotic administration, antibiotic treatment should not be delayed to facilitate this process.

Patients with Pontiac fever should not be prescribed antibiotic treatment. It is a self-limited illness that does not benefit from antibiotics, and patients usually recover within 1 week.

Treatment of extrapulmonary legionellosis varies according to the site of infection.

Reporting

Report all *Legionella* infections (or legionellosis cases) through the health department's web-based reportable disease surveillance system, PA-NEDSS (<https://www.nedss.state.pa.us/nedss/default.aspx>), or call the local health department or PADOH (877-PA-HEALTH). Timely reporting assists in the quick identification of potential clusters or outbreaks.

Any questions or concerns regarding these recommendations should be directed to the local health department or PADOH (877-PA-HEALTH).

Further information on legionellosis is available at:

Pennsylvania Department of Health: <https://www.health.pa.gov/topics/disease/Pages/Legionella.aspx>

Centers for Disease Control and Prevention (CDC): <https://www.cdc.gov/legionella/index.html>.

CDC's *Legionella* information for clinicians: <https://www.cdc.gov/legionella/hcp/clinical-guidance/>

Individuals interested in receiving future PA-HANs can register [here](#).

Categories of Health Alert messages:

Health Alert: conveys the highest level of importance; warrants immediate action or attention.

Health Advisory: provides important information for a specific incident or situation; may not require immediate action.

Health Update: provides updated information regarding an incident or situation; unlikely to require immediate action.

This information is current as of May 15, 2026, but may be modified in the future.
--