



A Publication of the Presbytery of West Virginia

520 2nd Ave, South Charleston, WV 25303 . wvpresbytery.org . 304-744-7634



Vol 4 Issue 10 ♦ October 2021

American Sleep Apnea Association
888-293-3650 (toll-free)

asaa@sleepapnea.org
www.sleepapnea.org

National Heart, Lung, and Blood Institute
301-592-8573
nhlbiinfo@nhlbi.nih.gov
www.nhlbi.nih.gov

National Sleep Foundation
703-243-1697
nsf@thensf.org
www.thensf.org
www.thensj.org/sleep-health-topics/

Restless Legs Syndrome Foundation
512-366-9109
info@rls.org
www.rls.org



Sleep and Aging (continued from September)

Sleep Apnea

People with sleep apnea have short pauses in breathing while they are asleep. These pauses may happen many times during the night. If not treated, sleep apnea can lead to other problems, such as high blood pressure, stroke, or memory loss.

You can have sleep apnea and not even know it. Feeling sleepy during the day and being told you are snoring loudly at night could be signs that you have sleep apnea.

If you think you have sleep apnea, see a doctor who can treat this sleep problem. You may need to learn to sleep in a position that keeps your airways open. Treatment using a continuous positive airway pressure (CPAP) device almost always helps people with sleep apnea. A dental device or surgery may also help.

Movement Disorders and Sleep

Restless legs syndrome, periodic limb movement disorder, and rapid eye movement sleep behavior disorder are common in older adults. These movement disorders can rob you of needed sleep.

People with restless legs syndrome, or RLS, feel like there is tingling, crawling, or pins and needles in one or both legs. This feeling is worse at night. See your doctor for more information about medicines to treat RLS.

Periodic limb movement disorder, or PLMD, causes people to jerk and kick their legs every 20 to 40 seconds during sleep. Medication, warm baths, exercise, and relaxation exercises can help.

Rapid eye movement, or REM, sleep behavior disorder is another condition that may make it harder to get a good night's sleep. Dur-

ing normal REM sleep, your muscles cannot move, so your body stays still. But, if you have REM sleep behavior disorder, your muscles can move and your sleep is disrupted.

Alzheimer's Disease and Sleep—A Special Problem

Alzheimer's disease often changes a person's sleeping habits. Some people with Alzheimer's disease sleep too much; others don't sleep enough. Some people wake up many times during the night; others wander or yell at night.

The person with Alzheimer's disease isn't the only one who loses sleep. Caregivers may have sleepless nights, leaving them tired for the challenges they face.


If you're caring for someone with Alzheimer's disease, take these steps to make him or her safer and help you sleep better at night:

- Make sure the floor is clear of objects.
- Lock up any medicines.
- Attach grab bars in the bathroom.
- Place a gate across the stairs.

Safe Sleep for Older Adults

Try to set up a safe and restful place to sleep. Make sure you have smoke alarms on each floor of your home. Before going to bed, lock all windows and doors that lead outside. Other ideas for a safe night's sleep are:

- Keep a telephone with emergency phone numbers by your bed.
- Have a lamp within reach that is easy to turn on.
- Put a glass of water next to the bed in case you wake up thirsty.
- Don't smoke, especially in bed.
- Remove area rugs so you won't trip if you get out of bed during the night.



Tips to Help You Fall Asleep

You may have heard about some tricks to help you fall asleep. You don't really have to count sheep—you could try counting slowly to 100. Some people find that playing mental games makes them sleepy. For example, tell yourself it is 5 minutes before you have to get up, and you're just trying to get a little bit more sleep.

Some people find that relaxing their bodies puts them to sleep. One way to do this is to imagine your toes are completely relaxed, then your feet, and then your ankles are completely relaxed. Work your way up the rest of your body, section by section. You may drift off to sleep before getting to the top of your head.

Use your bedroom only for sleeping. After turning off the light, give yourself about 20 minutes to fall asleep. If you're still awake and not drowsy, get out of bed. When you feel sleepy, go back to bed.

If you feel tired and unable to do your activities for more than 2 or 3 weeks, you may have a sleep problem. Talk with your doctor about changes you can make to get a better night's sleep.