

News | Articles | November 4, 2025

What Are the Infection Prevention Risks in the Hospital Tray Line?

Author(s) [Donald Sipp, MBA, RESE, CHESP, CHTI-2, CMIP, PMP](#)

"Hospital tray lines operate at high speed to feed hundreds of patients daily. Staff members plate food, cover trays, and load them onto mobile carts that navigate elevators, hallways, and patient rooms. Each of these steps introduces opportunities for contamination."



*Hospital cafeteria
(Gemeni AI-generated image
by author)*

Hospital meal trays are often overlooked in discussions of infection prevention. Yet each tray, from assembly to delivery, passes through multiple

hands, surfaces, and transport carts, transforming a routine process into a

potential vector for harmful pathogens.

While most attention in hospitals is focused



Advertisement

Advertisement

Trending on Infection Control Today

While most attention in hospitals is focused on medical equipment and patient care areas, the tray line, a seemingly simple part of the foodservice operation, represents a critical yet underappreciated pathway for infection transmission.

Hospital tray lines operate at high speed to feed hundreds of patients daily. Staff members plate food, cover trays, and load them onto mobile carts that navigate elevators, hallways, and patient rooms. Each of these steps introduces opportunities for contamination. Tray components, including lids, handles, utensils, and the carts themselves, can harbor microbes similar to those found on medical equipment, which studies indicate can persist even after standard cleaning procedures.^{1,2,3} This highlights the need to treat tray lines not simply as a food delivery mechanism but as a potential high-risk interface in the hospital environment.

ServSafe and Hospital Infection Control

Many hospitals rely on ServSafe certification to train foodservice staff in safe food handling, proper cooking temperatures, and the prevention of foodborne illnesses such as Salmonella and Listeria.⁴ ServSafe provides essential guidance for preventing traditional foodborne outbreaks, but it does

- 1 Why Infection Preventionists Belong in the C-Suite
- 2 "The Dirty Truth About Stethoscopes" and Other Must-Reads in ICT's November/December 2025 Print Issue
- 3 Oman's Clean Hospitals Initiative Sets a New Global Benchmark for Environmental Hygiene

not fully address health care-associated pathogens (HAPs), including methicillin-resistant *Staphylococcus aureus* (MRSA), vancomycin-resistant *Enterococci* (VRE), and *Clostridioides difficile* (*C difficile*). These pathogens can survive on high-touch surfaces, such as tray carts, lids, and utensils, posing an elevated risk to patients with weakened immune systems.^{2,3}

Integrating infection prevention principles beyond traditional food safety is crucial. Staff who understand environmental hygiene and proper handling techniques can dramatically reduce the potential for pathogen transfer along the tray line. Ensuring that foodservice training addresses both foodborne and HAPs strengthens the hospital's overall infection control strategy.

Evidence from Hospital Studies

Recent research underscores the risks associated with hospital foodservice operations. A 2025 study in Saudi Arabia documented significant microbial contamination throughout the hospital foodservice chain, affecting both raw ingredients and prepared meals.¹ Other studies examining hospital environmental surfaces have consistently shown persistent microbial presence despite standard cleaning protocols.^{2,3} While these studies

did not focus specifically on tray lines, they emphasize that mobile, high-touch surfaces remain potential reservoirs for infection. The cumulative effect of these surfaces, staff interactions, and patient contact points can significantly contribute to the overall risk of health care-associated infections.

Strategies for Safer Tray Lines

Hospitals can enhance food safety and reduce infection risks in tray lines through several targeted interventions:

- **Enhanced cleaning and disinfection** of tray carts, lids, and utensils to eliminate residual pathogens^{2,3}
- **Hand hygiene checkpoints** for staff handling trays, reinforcing proper techniques, and ensuring adherence
- **Dedicated transport routes** that separate foodservice operations from clinical areas to minimize cross-contamination
- **Integration of infection prevention training** into ServSafe or internal hospital certifications to address health care-specific pathogens⁴
- **Minimizing cross-touch points** through tray line design and workflow adjustments, such as automated or touchless systems

Even small operational changes, such as reorganizing tray line flow, staggering staff

reorganizing tray line flow, staggering staff shifts, or implementing routine disinfection checks, can significantly reduce infection risk.

Hospital tray lines are more than a logistical challenge; they are a frontline in infection prevention. While ServSafe establishes essential food safety standards,⁴ hospitals must supplement these with rigorous environmental hygiene, thoughtful workflow design, and ongoing staff education. By proactively addressing the unique risks associated with tray lines, health care facilities can better protect patients, limit the spread of pathogens, and strengthen the hospital's overall safety culture. Recognizing and mitigating the infection risks inherent in meal delivery is an important step toward a safer hospital environment for both patients and staff.

References

1. Alzahrani A, Al-Dawish M, Al-Mutairi A, et al. Comprehensive risk assessment and control measures in the food service chain of hospitals nutrition department: a case study in Al-Ahsa Governorate, Kingdom of Saudi Arabia. *Front Microbiol.* 2025;15:1551446. doi:10.3389/fmicb.2025.1551446
2. El-Sherbiny N, Khalil S, Hussein M, et al. The impact of enhanced cleaning on

bacterial contamination of hospital
environmental surfaces: a clinical trial in
critical care unit in an Egyptian hospital.
Antimicrob Resist Infect Control.
2024;13(22). doi:10.1186/s13756-024-
01489-z

3. Zhao Y, Liu W, Zhang X, et al.
Assessment of disinfection efficacy on
bacterial contamination of hospital
surfaces—China, 2007–2022. *China
CDC Wkly*. 2025;7(25):455-460.
doi:10.46234/ccdcw2025.135
4. National Restaurant Association.
ServSafe Manager Book. 8th ed.
Chicago, IL: National Restaurant
Association; 2023.

Newsletter

Stay prepared and protected with
Infection Control Today's
newsletter, delivering essential
updates, best practices, and expert
insights for infection preventionists.

Subscribe Now!

Related Content



Latest News: First Human Avian Influenza Case in 9...

By [Tori Whitacre Martonicz](#)

📅 November 14th 2025



Why Infection Preventionists Belong in the C-Suite

By

[Deborah Ellis, PhD, MS, MPH, MT\(ASCP\), CIC, LTC-CIP, CPHQ, FACHE](#)

📅 November 13th 2025



"The Dirty Truth About Stethoscopes" and...

By [Tori Whitacre Martonicz](#)

📅 November 12th 2025



Oman's Clean Hospitals Initiative Sets a New Global...

By

[Noel S. Gonzaga, MPH, BSMT, AMT, CIC, Jabir Al Sooti, CIC, CRCST](#)

📅 November 12th 2025



EPA Decision Opens New Chapter for Rechargeable...

By

Heather Stoltzfus, MPH, RN,
CIC,
Tori Whitacre Martonicz

📅 November 11th 2025

[Advertise](#)

[About Us](#)

[Editorial Board](#)

[Contact Us](#)

[Job Board](#)

[Terms and Conditions](#)

[Privacy](#)

[Do Not Sell My Personal Information](#)

[Cookie settings](#)

Contact Info



259 Prospect Plains Rd, Bldg H
Cranbury, NJ 08512



609-716-7777

MJH



© 2025 MJH Life Sciences®

All rights reserved.

[Home](#)

[About Us](#)

[News](#)

[Contact Us](#)