



Statin Therapy for Patients with Cardiovascular Disease – Received Statin Therapy

Measure Type: HEDIS (claims) / *Age Cohort*: 21-75 years (males); 40-75 years (females)

Measure Definition

Number of people who have had an MI, CABG, PCI, or other revascularization procedure* in the year prior or a diagnosis of ischemic vascular disease in both the measurement year and year prior and who were dispensed at least one high or moderate-intensity statin medication

*Endovascular Revascularization Lower Extremities

Documentation Required

- Diagnosis of cardiovascular disease:
 - Patient had MI (Inpatient Visit only) during year prior to measurement year
 - or
 - CABG, PCI, or other revascularization procedure during year prior to measurement year
 - or
 - Patient had ischemic vascular disease (IVD) during both measurement year and year prior (I20.0, I20.8, I20.9, I24.0, I24.8 - I25.119, I25.5 - I25.9, I63.20- I63.29, I63.50-I63.59, I65.01-I66.9, I67.2, I70.1-I70.799, I70.92, I75.011-I75.89, T82.855A, T82.855D, T82.855S, T82.856A, T82.856D, T82.856S)
- Dispensed at least one high or moderate-intensity statin medication ([Medication List F](#))

Applicable Visits: (Only one of the two visits may be a telehealth visit, a telephone visit or an online assessment).

- **CPT Codes:** 99201-99215, 99221-99233, 99238, 99239, 99241-99245, 99251-99255, 99291, 99341-99350, 99381-99397, 99401-99404, 99411, 99412, 99429, 99455, 99456
- **Telephone Visit CPT Codes:** 98966-98968, 99441-99443
- **Telehealth Modifier Codes:** GT, 95
- **Online Assessments Codes:** 98969, 99444

Note: Patients who meet any of the following criteria during measurement year or year prior are excluded from this metric: ESRD, cirrhosis, myalgia, myositis, myopathy, rhabdomyolysis, pregnancy, in vitro fertilization or were dispensed at least one prescription for clomiphene

Role of Care Management

- Ensure medication fill
- Reinforce patient education regarding chronic disease, medication adherence
- Identify & Address barriers to adherence

Role of PCP Practice

- Prescribe Statin Medication as appropriate
- Provide education to patient on diagnosis and symptom management



Statin Therapy for Patients with Cardiovascular Disease – Statin Adherence 80%

Measure Type: HEDIS (claims) / *Age Cohort*: 21-75 years (males); 40-75 years (females)

Measure Definition

Number of people who had an MI, CABG, PCI, or other revascularization procedure* in the year prior or a diagnosis of ischemic vascular disease (IVD) in both the measurement year and year prior and who achieved a proportion of days covered of 80% for the [treatment period](#)

*Endovascular Revascularization Lower Extremities

Documentation Required

- Diagnosis of cardiovascular disease:
 - Patient had MI (Inpatient Visit only) during year prior to measurement year
or
 - CABG, PCI, or other revascularization procedure during year prior to measurement year
or
 - Patient had ischemic vascular disease (IVD) during both measurement year and year prior (I20.0, I20.8, I20.9, I24.0, I24.8 - I25.119, I25.5 - I25.9, I63.20- I63.29, I63.50-I63.59, I65.01-I66.9, I67.2, I70.1-I70.799, I70.92, I75.011-I75.89, T82.855A, T82.855D, T82.855S, T82.856A, T82.856D, T82.856S)
- Dispensed at least one high or moderate-intensity statin medication ([Medication List F](#))
- Prescriptions are filled at least 80% of the time

Applicable Visits: (Only one of the two visits may be a telehealth visit, a telephone visit or an online assessment).

- **CPT Codes:** 99201-99215, 99221-99233, 99238, 99239, 99241-99245, 99251-99255, 99291, 99341-99350, 99381-99397, 99401-99404, 99411, 99412, 99429, 99455, 99456
- **Telephone Visit CPT Codes:** 98966-98968, 99441-99443
- **Telehealth Modifier Codes:** GT, 95
- **Online Assessments Codes:** 98969, 99444

Note: Patients who meet any of the following criteria during measurement year or year prior are excluded from this metric: ESRD, cirrhosis, myalgia, myositis, myopathy, rhabdomyolysis, pregnancy, in vitro fertilization or were dispensed at least one prescription for clomiphene

Role of Care Management

- Ensure monthly refills
- Reinforce patient education regarding chronic disease, medication adherence
- Identify & Address barriers to adherence

Role of PCP Practice

- Prescribe statin medication as appropriate and authorize monthly refills
- Provide education to patient on diagnosis and symptom management



Suffolk Care
Collaborative

MEASURE MEDICATION LISTS



Statin Therapy for Patients with Cardiovascular Disease – Received Statin Therapy & Statin Adherence 80%

[Back to Measure](#) - Received Statin Therapy

[Back to Measure](#) – Statin Adherence 80%



Description	Prescription
High-intensity statin therapy	<ul style="list-style-type: none">• Atorvastatin 40-80 mg• Amlodipine-atorvastatin 40-80 mg• Ezetimibe-atorvastatin 40-80 mg• Rosuvastatin 20-40 mg• Simvastatin 80 mg• Ezetimibe-simvastatin 80 mg
Moderate-intensity statin therapy	<ul style="list-style-type: none">• Atorvastatin 10-20 mg• Amlodipine-atorvastatin 10-20 mg• Ezetimibe-atorvastatin 10-20 mg• Rosuvastatin 5-10 mg• Simvastatin 20-40 mg• Ezetimibe-simvastatin 20-40 mg• Niacin-simvastatin 20-40 mg• Sitagliptin-simvastatin 20-40 mg• Pravastatin 40-80 mg• Lovastatin 40 mg• Niacin-lovastatin 40 mg• Fluvastatin XL 80 mg• Fluvastatin 40 mg bid• Pitavastatin 2–4 mg