



# AMERICAN SOCIETY OF INTERVENTIONAL PAIN PHYSICIANS

THE VOICE OF INTERVENTIONAL PAIN MANAGEMENT

## ASIPP Risk Stratification of Patients Presenting for Interventional Pain Procedures: Decreasing Morbidity of COVID-19

If Patient Residence status is **Nursing Home or Assisted Living Facility or Incarceration** during the past 30 days, consider as **HIGH-RISK** Patient. If not follow below table to Risk Stratification table.

**Risk Stratification Table**

RISK FACTOR	LOW RISK	MODERATE RISK	HIGH RISK	COMMENTS
Age (years)	45-64 years (1)	65-74 years (2)	>= 75 years (3)	
Pulmonary	None (0)	Mild Intermittent Asthma (2)	Chronic Lung Condition i.e. Moderate to Severe Asthma, COPD (3)	
Cardiovascular	None (0)	HTN or CAD (2)	HTN + CAD HTN + CHF HTN + CAD + CHF CHF alone (3)	
Obesity	BMI 24.9-29.9 (1)	BMI 30.0-39.9 (2)	BMI >= 40 (3)	
Diabetes (A1C) (BGM mg/dl)	5.7-6.49 or 100-120 mg/dl (1)	6.50-8.49 or 120-160 mg/dl (2)	>= 8.5 or >160 mg/dl (3)	Consider finger stick BGM if A1C not available
Renal	None (0)	Acute or Chronic Renal Insufficiency (2)	Chronic Renal Insufficiency on Dialysis (3)	
Hepatic	None (0)	Chronic Hepatitis (2)	Cirrhosis (3)	
Immuno- Compromised State	None (0)	One stable condition (2)	Present of <b>ANY</b> -Cancer (active treatment) -BM/Organ Transplantation -Immune Deficiencies -Poorly Controlled HIV/AIDS -Chronic Steroid Use (3)	

**\*\* The scoring number for risk factor is listed in parentheses. This number indicates the points for that condition.**

**Low Risk: <= 7 points**

**Moderate Risk: 8-14 points**

**High Risk: >= 15 points**

ASIPP Return to Practice Task Force Members: Shalini Shah (Chair) , Sudhir Diwan, Amol Soin, Christopher Gharibo, Kartic Rajput