

Appendix K.

Guidelines for Young Adult Offenders

Adopted February 10, 2017

I. Introduction

The purpose of this appendix is to provide Multi-Disciplinary Treatment Teams (MTTs) with additional guidance on working with domestic violence offenders ages 18-25, who can be classified as young adults (note, this population is also sometimes referred to as transitioned-aged). This informational document provides MTTs with best practices guidelines, potential risk and protective factors, and suggestions for the treatment and case management of young adults. The guidelines in this appendix do not replace any of the mandates currently required in the *Standards*. A Young Adult Committee of the DVOMB was convened to develop these Guidelines. The Committee, comprised of state and local experts in the field of at-risk youth, delinquent juveniles, and young adult populations (including treatment providers, victim service providers and advocates, probation/corrections officers, and others involved in the criminal justice system) collaborated on the creation of this Appendix. Clinical and professional expertise, as well as a review of available research and literature, served as the foundation for these Guidelines.

The Risk, Need, and Responsivity (RNR) principles are an evidence-based framework for evaluating, treating and supervising individuals involved with the criminal justice system. The RNR principles originated from numerous high-quality and generalizable studies in the broader criminological literature (Andrews & Bonta, 2010). The RNR principles state the following:

- Risk - Services provided to offenders should be proportionate to the offenders' relative level of static and dynamic risk (i.e., low, moderate, or high risk) based upon accurate and valid research-supported risk assessment instruments;
- Need - Interventions are most effective if services target criminogenic needs (both social and psychological factors) that have been empirically associated with recidivism; and
- Responsivity - Effective service delivery of treatment and supervision requires individualization that matches the offender's strengths, culture, learning style, and abilities, among other factors. (Please see Appendix E (VI) - Responsivity Principle and Factors).

The DVOMB recognizes that based on responsivity issues and the needs of young adults, a different approach may be needed when addressing the unique challenges of this population. Neurobiological research gives us a deeper understanding of adolescent, young adult brain

Colorado Domestic Violence Offender Management Board

700 Kipling Street, Suite 3000

Denver, Colorado 80215

development, and neuro-psychology.¹ This research indicates that the brains of young adults are more fluid, and are still developing and changing until the age of 25 (Perry, 2009; Spear, 2010, Teicher, 2002). As a result, some young adults may not recognize the consequences of their behaviors and may present more like an adolescent rather than an adult. Research indicates that over-responding to non-criminal violations with this population can cause more harm than good for the offender, victim and community (Teicher, 2002). As a result, young adults are at higher risk for dropping out of treatment (Buttell & Carney, 2008; Jewell & Wormith, 2010), therefore, it is imperative for MTT members to assess and treat this population within a framework that is appropriate for this population.

Disclaimer of Risk and Safety Issues: The information outlined in this Appendix does not replace, change or supersede the risk factors identified by the Domestic Violence Risk and Needs Assessment (DVRNA) as part of the Offender Evaluation. These guidelines offer recommendations to lower risk and enhance responsibility by increasing treatment readiness and amenability to make positive behavioral change. Additional offender competencies are necessary to address specific issues unique to the development of the young adult population that may not be currently addressed by Standard 5.08. Appropriate interventions should be commensurate with the nature and severity of the behavior and the degree to which it relates to risk. Risk of harm to others must not be ignored and should be balanced when assessing impulsive behavior typical in adolescence versus more criminal and anti-social characteristics, which are indicative of heightened risk.

II. Guiding Principles

The Guiding Principles (described in Section 3.0) are designed to assist and guide the work of those involved in the management and containment of domestic violence offenders. For the purposes of this Appendix, the following guiding principles in the Standards may or may not be relevant or appropriate for young adult offenders.

Guiding Principle	Issue for Consideration with Young Adults
GP 3.06 - It is the nature of domestic violence offenders that their behaviors tend to be covert, deceptive, and secretive. These behaviors are often present long before they are recognized publicly.	Young adults may not have a prior history of domestic violence due to their age, and a history of delinquency may be more prevalent. Impulsivity and poor decision-making may be attributed to the index offense. As a result, deceptiveness and secrecy may not be as normative with young adults.

¹ During adolescence, the human brain experiences increased growth, connectivity, and synaptic pruning (Spear, 2010). The rate at which the development of the neural pathways associated with regulation and reward sensitivity may provide insight into the characteristics of emerging adulthood.

Colorado Domestic Violence Offender Management Board

700 Kipling Street, Suite 3000

Denver, Colorado 80215

GP 3.13 - The preferred treatment modality is group therapy.	Research and clinical experience indicates that young adults tend to respond better to a combination of both individual and group sessions. If not, negative therapeutic outcomes may occur by exposing young adults to older adults who are higher risk and developmentally more mature (Abracen et al., 2016; Lowenkamp & Latessa, 2004).
GP 3.15 - Offender treatment must address the full spectrum of abusive and controlling behaviors associated with domestic violence, and not just the legally defined criminal behavior(s).	Young adults may be at a developmental stage where maladaptive patterns of violence (with intimate partners and significant others) may not fully be present, but may or may not require an intervention that addresses the full-spectrum of abusive and controlling behaviors.

III. Young Adult Risk and Protective Factor Considerations

Given the emerging research on young adults, it is important for the MTT to evaluate an offender's problematic behavior and assess their developmental maturation. When responding to rule breaking or non-compliance, it is best to determine whether or not it signifies an increase in risk. If so, the MTT should assess what needs exist and what intervention best addresses those needs and manages risks appropriately. Such assessment should include strengths and protective factors.²

Contributing risk factors in young adults will likely be best mitigated by ensuring the MTT prioritizes the RNR principles and ensures all of these are assessed and addressed in treatment. The DVRNA is based on adult risk factors and does not address protective factors that may be present within young adult clients. Providers should consider research-supported developmentally-appropriate risk and protective factors in the ongoing assessment and case management of young adults (see attachment 1 for examples of potential risk and protective factors for this population). Providers should exercise clinical judgement with these cases in terms of identification of risk and protective factors, and consult with other professionals as needed.

Risk Assessment Instruments and Collateral Information

The DVRNA instrument still applies to the young adult population. However, young adults may possess issues that require further assessment and consideration of risk and needs. Adult risk

² Protective factors are conditions or attributes (skills, strengths, resources, supports or coping strategies) in individuals, families, communities or the larger society that help people deal more effectively with stressful events and mitigate or eliminate risk in families and communities.

Colorado Domestic Violence Offender Management Board

700 Kipling Street, Suite 3000

Denver, Colorado 80215

assessment instruments may or may not necessarily address the unique factors of the young adult population. After conducting a thorough evaluation, in accordance with Section 4.0 of the Standards, alternative risk assessments may be appropriate in some cases to use for juveniles under the age of 18 for informational purposes only. It is important to note that using a juvenile risk assessment instrument on an individual over the age of 18 is not a validated assessment of risk, but may be informative for case planning.

IV. Responsivity Issues Affecting Young Adults to Consider

The MTT should individualize treatment and supervision for young adults, to the extent possible, in relation to their present development, deficits and amenability to treatment. Group-specific sessions for young adults should be considered and utilized by providers (where applicable), in order to minimize the exposure of young adults to older adults who are higher risk and developmentally more mature. Additionally, it may be more appropriate for young adults to receive individual sessions instead of group sessions based on the clinical judgement of a provider. Providers utilizing individual sessions with young adults should develop treatment plans that address the client's individual needs, their Core Competencies and other unique issues present.

It is important for the MTT to also understand the characteristics of offenders that may be more likely to lead to treatment dropout. The research indicates that a client is more likely to complete treatment if they are older and employed (Buttell & Carney, 2008; Jewell & Wormith, 2010). As a result, the challenges associated with emerging adulthood such as finding stable housing, employment and relationships may increase treatment dropout for young adults. Providers should consider any healthy and pro-social factors that may assist a young adult client's initial engagement in the therapeutic process and commitment over time, and focus on development of protective factors.

V. MTT Guidelines for Decision-Making

MTTs are encouraged to assess and develop individualized treatment plans including containment efforts for young adults, based on their maturation and risk levels. Independent living skills, risk, and protective factors should be discussed by MTTs and factored into programming for the offender. MTTs should consult with other experienced adult and juvenile practitioners to assist in the development of effective treatment and supervision, and to identify possible resources that may aid in information gathering, where such experience is lacking.

Recommendations:

- Use an evidence-based, research-informed or best practice curriculum for this population (see for example, Gibbes, L. & Myers, L., 2011)
- Consider the following treatment issues, among others:

Colorado Domestic Violence Offender Management Board

700 Kipling Street, Suite 3000

Denver, Colorado 80215

- Self-efficacy and self-identity
- Empathy
- Developmental stage
- Appropriate boundaries and communication in relationship
- Age appropriate healthy sexuality
- Age appropriate healthy sexual behavior
- Appropriate use of electronic devices and social media
- Appropriate dating skills
- Positive support groups and peer associations
- Consider additional core competencies, as appropriate (See Section 5.08, VI)
- Support ongoing research to better inform interventions with this population
- Stay current on research related to this population, including developmental issues
- Consult with other professionals as needed
- Participate in trainings on:
 - Human development and maturation of young adults
 - Brain development and neuro-psychology

VI. Links to Resource Documents

Gibbes, L. & Myers, L. (2011). Colorado Teen Dating Violence Prevention Final Report Primary Prevention of Teen Dating Violence in the Denver-Aurora Community: Best Practices and Strategy Recommendations. *Colorado Department of Public Health and Environment*.

[Click Here](#)

Oudekerk, B., Blachman-Demner, D., & Mulford, C. (2014). Teen Dating Violence: How Peers Can Affect Risk & Protective Factors. *U.S. Department of Justice, Office of Justice Programs, National Institute of Justice, NCJ 248337*.

[Click Here](#)

Colorado Domestic Violence Offender Management Board

700 Kipling Street, Suite 3000

Denver, Colorado 80215

Attachment 1 - Examples of Potential Risk and Protective Factors for this Population

This table is meant to be a resource that lists some possible risk and protective factors associated with youth that may be considered for young adults. This is not exhaustive list of risk or protective factors as there may be others that should be considered by the MTT. Providers should consider research-supported developmentally-appropriate risk and protective factors in the ongoing assessment and case management of young adults. Providers should exercise clinical judgement with these cases in terms of identification of risk and protective factors, and consult with other professionals as needed.

Table of Risk and Protective Factor Chart

Domains	Risk Factors	Adolescent Problem Behaviors					Protective Factors
		Substance Use	Depression/ Anxiety	Delinquency	Teen Pregnancy	School Dropout	
	Risk factors increase the likelihood youth will develop problem behaviors						Protective factors help protect or buffer the risks of youth developing problem behaviors.
Community	Availability of alcohol/other drugs	X					X
	Availability of firearms		X				X
	Community laws and norms are favorable toward drug use, firearms and crime	X	X				X
	Transitions and mobility	X	X	X		X	
	Low neighborhood attachment and community disorganization	X		X			X
	Media portrayals of violence	X		X			X
	Extreme economic deprivation	X		X	X	X	X
Family	Family history of problem behavior	X	X	X	X	X	X
	Family management problems	X	X	X	X	X	X
	Family conflict	X	X	X	X	X	X
	Favorable parental attitudes	X		X			X
School	Academic failure beginning in late elementary school	X		X	X	X	X
	Lack of commitment to school	X		X	X	X	X
Individual/ Peer	Early and persistent antisocial behavior	X		X	X	X	X
	Rebelliousness	X	X	X		X	
	Friends who engage in the problem behavior	X		X	X	X	X
	Favorable attitudes toward the problem behavior	X		X	X	X	
	Early initiation of the problem behavior	X		X	X	X	X
	Gang involvement	X		X			X
	Constitutional factors	X	X	X			X

Source: Hawkins, J., Catalano, R., Arthur, M. (2002). Promoting science-based prevention in communities. *Addictive Behavior*, 27(6):951-76.

Colorado Domestic Violence Offender Management Board

700 Kipling Street, Suite 3000

Denver, Colorado 80215

References

Abracen, J., Gallo, A., Looman, J., & Goodwill, A. (2015). Individual community-based treatment of offenders with mental illness: relationship to recidivism. *Journal of Interpersonal Violence*, 1- 17.

Andrews, D., & Bonta, J. (2010). The psychology of criminal conduct (5th ed.). New Providence, NJ: LexisNexis Matthew Bender.

Buttell, F., & Carney, M. (2008). A Large Sample Investigation of Batterer Intervention Program Attrition: Evaluating the Impact of State Program Standards. *Research on Social Work Practice*, 18(3):177-188.

Gibbes, L. & Myers, L. (2011). Colorado Teen Dating Violence Prevention Final Report Primary Prevention of Teen Dating Violence in the Denver-Aurora Community: Best Practices and Strategy Recommendations. *Colorado Department of Public Health and Environment*.

Gover, A., Jennings, W., Davis, C., Tomisch, E., and Tewksbury, R. (2011). Factors related to the completion of domestic violence offender treatment: The Colorado experience. *Victims and Offenders*, 137-156.

Hawkins, J., Catalano, R., Arthur, M. (2002). Promoting science-based prevention in communities. *Addictive Behavior*, 27(6):951-76.

Jewell, L., & Wormith, S. (2010). Variables Associated with Attrition from Domestic Violence Treatment Programs Targeting Male Batters: A Meta-Analysis. *Criminal Justice and Behavior*. 37(10):1086-1113.

Lowenkamp, C., & Latessa, E. (2004). Increasing the effectiveness of correctional programming through the risk principle: Identifying offenders for residential placement. *Criminology and Public Policy*, 4, 501– 528.

Oudekerk, B., Blachman-Demner, D., & Mulford, C. (2014). Teen Dating Violence: How Peers Can Affect Risk & Protective Factors. *U.S. Department of Justice, Office of Justice Programs, National Institute of Justice, NCJ 248337*.

Perry, B. (2009). Examining Child Maltreatment Through a Neurodevelopmental Lens: Clinical Applications of the Neurosequential Model of Therapeutics. *Journal of Loss and Trauma*, 14:240–255.

Spear, L. (2010). The behavioral neuroscience of adolescence. New York: W.W. Norton & Company Inc

Colorado Domestic Violence Offender Management Board

700 Kipling Street, Suite 3000

Denver, Colorado 80215

Teicher, M., Anderson, S., Polcari, A., Anderson, C., & Navalta, C. (2002). Developmental neurobiology of childhood stress and trauma. *Psychiatric Clinics of North America*, 25, 397-426.