

WuXi *In vitro* Immunology Assay Platform



WuXi AppTec Research Service Division, Oncology & Immunology Unit





2019.09

Service list of *In-vitro* and *ex-vivo* immunology assays


Assay			WuXi capability (Human/Mouse)
Cell Activation		T Cell Agonism (Antibody/SEB/CMV Recall)	Yes (H/M)
		B Cell Activation	Yes (H/M)
		NK Cell Activation/Expansion	Yes (H/M)
Cell Differentiation	T Cell Polarization	Treg	Yes (H/M)
		TH1	Yes (H/M)
		TH2	Yes (H/M)
		TH17	Yes (H/M)
		Macrophage Differentiation	
		M1/M2 Macrophage	Yes (H/M)
		MDSC	Yes (H)
		DC/Tumor activate DC	Yes (H/M)
Cytotoxicity Assay	T Cell	OT-1 system	Yes (M)
	NK Cell	NK cell cytotoxicity	Yes (H)
		ADCC/CDC/ADCP	Yes (H)
Mixed Lymphocyte Reaction		DC co-culture with T	Yes (H/M)
Immune Cell Suppressive Assay		MDSC suppressive assay	Yes (H)
		Treg suppressive assay	Yes (H)
		Tumor cell suppressive assay (co-culture with immune cells)	Yes (H)
		Sting pathway activation	
Other assay		NLRP3 inflammasome activation assay	Yes (H)

T Cell Activation by Antibody

Anti-CD3/CD28 Stimulation



T Cell Activation by SEB








CMV Recall Assay

CD4+ T


Before stimulation

After stimulation




B Cell Activation

Anti-IgM Stimulation




Primary NK Activation




Primary NK Expansion

No
stimulus



stimulation



T helper Cell Polarization

Treg differentiation

Day 0

Isolation of naïve CD4⁺ T cells from PBMC




Stimulation with anti-CD3/CD28 in the presence of
Treg differentiation cocktail



Day 5

FACS (LIVE-DEAD/CD4/CD25/FoxP3)



T helper Cell Polarization

Th1 differentiation

Day 0

Isolation of naïve CD4⁺ T cells from PBMC




Stimulation with anti-CD3/CD28 Dynabeads in the presence of Th1 differentiation cocktail



Day 5

FACS (LIVE-DEAD/CD4/IFN- γ)



T helper Cell Polarization

Th2 differentiation

Day 0

Isolation of naïve CD4⁺ T cells from PBMC




Stimulation with anti-CD3/CD28 in the presence of
Th2 differentiation cocktail



Day 13

FACS (LIVE-DEAD/CD4/IL-4)



T helper Cell Polarization

Th17 differentiation

Day 0

Isolation of naïve CD4⁺ T cells from PBMC




Stimulation with anti-CD3/CD28 in the presence of
Th17 differentiation cocktail





Day 11

FACS (LIVE-DEAD/CD4/IL-17A)

Th0 condition




Th17 condition





Isotype

Macrophage Differentiation

Human M1/M2 macrophage differentiation





FACS analysis of M1/M2 macrophage:



CTL Cytotoxicity Assay (OT-I System)


- **Effector:** Antigen-specific OT-I splenocyte culture
- **Target:** CellTrace Far-Red labeled EL4 cells




NK92MI Cyto-K562

K562 cells

K562 only




FSC-A



PI


CellTrace




PI

CellTrace

NK92MI+K562




FSC-A



PI

CellTrace





PI

CellTrace

NK Cytotoxicity Assay (NK-92MI Cell Line)

- **Effector:** NK-92MI cells
- **Target:** K562 cells labeled with 100 nM CellTrace Far-Red



Duplicate wells of E/T ratios:




ADCC Assay (Purified Primary NK cells)

Antibody-Dependent Cell-Mediated Cytotoxicity

- **Effector:** NK cells isolated from PBMC of healthy donor
- **Target:** CD20⁺ Raji cell line, labeled with 50 nM CellTrace Far-Red





ADCP Assay-M-CSF induced macrophage





Mixed Lymphocyte Reaction

DC co-cultured with T cell



Mixed Lymphocyte Reaction


DC co-cultured with T cell





MDSC Suppressive Assay

MDSC Induction and Co-culture with T Cell

Induction of MDSC from PBMC





MDSC suppress T cell proliferation and cytokine production in co-culture



Treg Suppressive Assay



CO-culture with Teff



Tumor Cell Suppressive Assay


Co-culture with Immune Cells

- Hela cell co-cultured with primary human T cells in the presence of cpdX.




STING Pathway Activation

THP-1 (PMA)




PBMC



NLRP3 Inflammasome Activation

- PMA-differentiated THP-1 cells were stimulated with LPS, followed by treatment with Nigericin for 1h





OUR COMMITMENT

Improving Health. Making a Difference.

For questions and requests, please email to info_onco@wuxiapptec.com



<https://onco.wuxiapptec.com>



Mobile App

 药明康德
WuXi AppTec
www.wuxiapptec.com