

OTOTOXICITY RISK AWARENESS




HEARING HEALTH DECISIONS

IN THE U.S., OVER 100 CLASSES OF DRUGS COMMONLY USED TO TREAT PAIN AND INFECTIONS, AS WELL AS CANCER, HEART OR KIDNEY DISEASE, CAN **DAMAGE THE INNER EAR.**^[1]

DID YOU KNOW?

The manner in which certain drugs can cause ringing in the ears (Tinnitus),^[2] followed by hearing loss or dizziness, is known as “ototoxicity.”




TALK TO YOUR CARING DOCTORS:

Who will help educate you about the hearing health-related side effects of treatment-related medications.

OUR EXPERT ADVICE CAN HELP!

GET EXPERT HEARING TESTING:

Including baseline and regular screenings which will enable early detection of ototoxic impact as well as timely response if and when hearing loss occurs.



©2016 HEALTHSCAPES®

[1] American Academy of Audiology Position Statement and Clinical Practice Guidelines Ototoxicity Monitoring. audiology.org 2009 [2] Ototoxic Medications, Drugs that can cause hearing loss and tinnitus. League for the Hard of Hearing 2000 [3] Schacht, *et al*: Cisplatin and Aminoglycoside Antibiotics: Hearing Loss and Its Prevention, *The Anatomical Record* (Hoboken). 2012 [4] Rybak and Ramkumar, Ototoxicity. *Kidney International* 2007