Hidden Hormone Disruptors: Toxins, Hormonal Balance & Fertility

A naturopathic guide for women and men

November 2026

Contents

Introduction 3
Chapter 1: How hormones control fertility 5
Chapter 2: What endocrine disruptors are and how they act 6
Chapter 3: Plastics chemicals — bisphenols and phthalates 8
Chapter 4: “Forever chemicals” — PFAS 10
Chapter 5: Pesticides and herbicides 12
Chapter 6: Personal care and household chemicals 14
Chapter 7: Heavy metals, air pollution and lifestyle toxins 16
Chapter 8: Effects on women’s fertility 18
Chapter 9: Effects on men’s fertility 20
Chapter 10: Your practical action plan 22
Chapter 11: Lowering your toxic burden naturally 24
Chapter 12: Testing, your practitioners, and keeping perspective 28
Quick reference 30
Glossary 32
Sources 33

Introduction

Everyday chemicals that interfere with hormones — called endocrine-disrupting chemicals, or EDCs — are now consistently linked to poorer fertility in both women and men. A 2025 systematic review of a decade of human studies found consistent associations between EDC exposure and impaired semen quality, lower ovarian reserve, infertility, PCOS and altered hormone levels.

The good news: many of the biggest exposures come from food, packaging, water, cosmetics and household products — things you can change. This e-book explains what these chemicals are, where they hide, what the evidence actually shows, and a realistic plan for lowering your load.

A naturopathic approach. This e-book follows the core principles of naturopathy, applied with the best available science:

- **Identify and remove the cause** - reduce exposures at their source before anything else.
- **Support the healing power of nature** - your liver, gut, kidneys, skin and lungs already clear toxins every day; good nutrition, movement, hydration, sleep and calm help them do it well.
- **First, do no harm** - choose the gentlest effective steps and avoid harsh or unproven "detox" regimes.
- **Treat the whole person** - hormones respond to diet, stress, sleep, relationships and environment together.
- **Prevention** - the months before conception are a powerful window to build health in both partners.
- **Doctor as teacher** - understanding where toxins come from lets you make informed choices.

How to use this e-book

- Chapters 1–2 explain how hormones control fertility and how chemicals interfere with them.
- Chapters 3–7 profile the main chemical families: where they come from and what they do.
- Chapters 8–9 summarize effects on women and on men.
- Chapter 10 is a step-by-step plan for reducing exposure.
- Chapter 11 shows how nutrition, physical activity, hydration, sleep and stress management support your body's natural elimination pathways.
- Chapter 12 covers testing and working with your naturopathic and medical practitioners.
- The final section has quick-reference tables, a checklist, a glossary and sources.

A note on the evidence. Most human studies are observational: they show that higher exposure tends to go with worse outcomes, not that a chemical definitely caused a specific person's infertility. Infertility usually has several causes at once. Use this information to reduce avoidable risks, not to blame yourself.

Medical disclaimer. This e-book is general education, not medical advice. It works best alongside care from a licensed naturopathic doctor and your medical team. If you have been trying to conceive for 12 months (6 months if you are 35 or older), have irregular cycles, or have known reproductive conditions, see a doctor or fertility specialist.

Chapter 1: How hormones control fertility

Fertility in both sexes runs on one chain of command — the hypothalamic–pituitary–gonadal (HPG) axis — and small signals at any link can shift the whole system. That is why chemicals active at tiny doses can matter.

The chain of command

1. **Hypothalamus (brain)** releases GnRH in pulses.
2. **Pituitary gland** responds by releasing two messengers: FSH (follicle-stimulating hormone) and LH (luteinizing hormone).
3. **Gonads** — ovaries or testes — respond by maturing eggs or sperm and making sex hormones.
4. **Feedback**: sex hormones travel back to the brain, which turns GnRH, FSH and LH up or down to keep levels balanced.

In women

- FSH grows a follicle each cycle; the follicle makes **estrogen**, which builds the uterine lining.
- An LH surge triggers ovulation. The empty follicle (corpus luteum) then makes **progesterone** to support a possible pregnancy.
- **AMH** (anti-Müllerian hormone) reflects the remaining egg supply, or ovarian reserve.

In men

- LH tells Leydig cells in the testes to make **testosterone**.
- FSH and testosterone together drive sperm production in Sertoli cells. A full cycle of sperm production takes roughly 70–90 days — so changes you make today show up in semen about three months later.

The supporting cast

- **Thyroid hormones** set metabolic pace; both underactive and overactive thyroid disrupt cycles and sperm.
- **Insulin** matters too: insulin resistance drives excess androgens in PCOS.
- **Cortisol** (stress hormone) can suppress the HPG axis.

Endocrine disruptors can act at every level of this chain — brain, pituitary, gonads, thyroid and metabolism — which is why their effects are so varied.

Chapter 2: What endocrine disruptors are and how they act

An endocrine disruptor is any outside chemical that interferes with how your body makes, moves, uses or clears hormones. Researchers describe EDCs as acting by mimicking, blocking, or altering the synthesis, transport, metabolism or elimination of natural hormones.

Five ways they interfere

- **Mimic** — fit into a hormone receptor and switch it on. BPA, for example, weakly imitates estrogen.
- **Block** — sit in a receptor without switching it on, so the real hormone can't act. Many phthalates and some pesticides are anti-androgenic (they oppose testosterone).
- **Change production** — interfere with the enzymes that build hormones, such as steroid production in Leydig cells or ovarian follicles.
- **Change transport and clearance** — alter the proteins that carry hormones in blood, or how fast the liver breaks them down.
- **Damage cells directly** — trigger oxidative stress and inflammation, which harm eggs, sperm and sperm DNA.

Why "the dose makes the poison" doesn't fully apply

- **Low doses can matter.** Hormones work at parts-per-billion levels, so tiny amounts of a mimic can tip the balance. Some EDCs show non-linear effects, where a low dose acts differently from a high one.
- **Timing matters most.** Exposure in the womb, in early childhood and at puberty can program the reproductive system for life. A boy's exposure before birth may shape his sperm count decades later.
- **Mixtures add up.** We are exposed to dozens of chemicals at once. Several weak anti-androgens together can have a combined effect, even if each alone is below a "safe" level.
- **Persistence varies.** Some chemicals leave the body within hours to days (BPA, phthalates, parabens), so cutting exposure lowers levels fast. Others stay for years (PFAS, PCBs, DDT, lead in bone), so prevention is the main tool.

How they get in

- **Eating and drinking** — the largest route for most people: packaging, can linings, processed food, pesticide residues, contaminated water, fatty animal foods for persistent chemicals.
- **Skin** — lotions, cosmetics, fragrance, sunscreen, receipts.

- **Breathing** — household dust, air fresheners, vehicle and industrial pollution, smoking.
- **Work** — farming, manufacturing, salons, painting, firefighting and plastics work can bring much higher exposures.

Chapter 3: Plastics chemicals — bisphenols and phthalates

Bisphenols and phthalates are the most common hormone disruptors in daily life, and also the easiest to cut: both leave the body within hours to days. In a 2011 study, families who ate only fresh, unpackaged food for three days saw urine BPA fall by 66% and DEHP metabolites by 53–56%.

Bisphenol A (BPA) and its substitutes

What it is. A building block of hard polycarbonate plastic and the epoxy linings of food and drink cans.

Where you meet it

- Canned food and drinks (can linings), especially acidic foods
- Hard reusable plastic bottles and containers (recycling code 7, "PC")
- Thermal paper receipts
- Some dental sealants and older water pipe coatings

How it acts. BPA weakly mimics estrogen and can interfere with androgen and thyroid signaling. A 2025 review called BPA and the phthalate DEHP among the most widespread EDCs affecting female fertility.

"BPA-free" caveat. Many products swapped BPA for BPS or BPF, which act similarly in lab studies. Treat "BPA-free" as "bisphenol-different," not "hormone-safe."

Regulation. The EU banned BPA and other hazardous bisphenols in food-contact materials under Regulation (EU) 2024/3190, effective 20 January 2025, with transition periods. This followed a 2023 EFSA reassessment concluding that typical BPA exposure posed a health risk. In the US, BPA remains permitted in most food packaging.

Phthalates

What they are. "Plasticizers" that make plastic soft and flexible, and solvents that help scents last.

Where you meet them

- Soft PVC/vinyl (recycling code 3): shower curtains, flooring, tubing, some food wrap
- Processed and fast food — they leach from plastic tubing and gloves in food production
- "Fragrance" or "parfum" in cosmetics, lotions, cleaners and air fresheners (phthalates are rarely listed by name)
- Nail polish and some hair products

- Household dust

How they act. Several phthalates are anti-androgenic: they lower testosterone production. A systematic review for the US EPA found robust evidence linking DEHP and DBP to male reproductive outcomes such as semen quality, testosterone and anogenital distance, and moderate evidence for DINP and BBP. In women, phthalates have been linked to endometriosis, fewer mature eggs and poorer IVF outcomes.

Key names to know: DEHP, DBP (DnBP), DiBP, BBP, DINP, DEP.

What to do

- Choose fresh or frozen food over canned; choose glass, stainless steel or cardboard cartons.
- ***Never*** heat food in plastic - no microwaving in plastic containers, no hot liquids into plastic.
- Cook at home more; fast food and takeout raise phthalate levels.
- Decline paper receipts, or handle them briefly and wash hands (especially before eating).
- Pick fragrance-free personal care and cleaning products.
- Avoid vinyl (PVC, code 3) where you can; swap plastic shower curtains for fabric.

Chapter 4: “Forever chemicals” — PFAS

PFAS stay in the body for years, so the strategy is to stop new intake, especially water and food packaging. In a 2023 Singapore study of women trying to conceive, those with higher blood PFAS had roughly a 30–40% lower chance of pregnancy or live birth within a year.

What they are. Per- and polyfluoroalkyl substances: thousands of man-made chemicals that repel water, grease and stains. Their carbon–fluorine bond almost never breaks down, so they build up in the environment and in people. PFOA and PFOS have half-lives in the human body of several years.

Where you meet them

- Contaminated drinking water — near airports, military bases, factories and sites using firefighting foam
- Grease-resistant food packaging: microwave popcorn bags, fast-food wrappers, pizza boxes, some takeout containers
- Older or scratched non-stick cookware
- Stain- and water-resistant carpets, upholstery, outdoor clothing
- Some cosmetics (look for “PTFE” or “perfluoro-” ingredients), dental floss and ski wax
- Fish from contaminated waters

Effects on women

- **Fertility:** in the Singapore preconception study of 382 women, each step up in exposure to individual PFAS cut fecundability by 5–10%. PFDA, PFOS, PFOA and PFHpA contributed most.
- **PCOS and endometriosis:** women with PCOS tend to have higher PFAS levels, and PFAS have been linked to endometriosis and delayed puberty.
- **Pregnancy:** PFAS cross the placenta and have been linked to lower birth weight and pregnancy-related high blood pressure.

Effects on men

- A study of 105 young Danish men found that those with high combined PFOS and PFOA had a median of 6.2 million normal sperm vs 15.5 million among men with low levels.
- Lab studies suggest PFAS can interfere with testosterone production and damage the testes’ blood barrier.

Regulation (as of late 2026). In the US, EPA’s 2024 rule set enforceable drinking-water limits of 4 parts per trillion for PFOA and PFOS. Those limits are being kept, but on 18 May 2026 EPA proposed rescinding limits for four other PFAS and extending compliance deadlines. Several US states keep their own stricter rules, and the EU is moving to restrict PFAS more broadly.

What to do

- Check your water utility's annual Consumer Confidence Report, or test a private well.
- If PFAS are present, use a filter certified to NSF/ANSI 53 or 58 for PFOA/PFOS. Reverse osmosis and good activated-carbon filters both reduce PFAS.
- Replace scratched non-stick pans with stainless steel, cast iron, carbon steel or ceramic-coated cookware.
- Pop popcorn on the stove; eat less fast food and greasy takeout in coated wrappers.
- Skip stain-guard treatments on new furniture and carpets.
- Dust with a damp cloth and vacuum with a HEPA filter — PFAS collect in household dust.

Chapter 5: Pesticides and herbicides

The goal is not to eat less produce but to eat less pesticide. In Harvard's EARTH study, men eating 1.5+ daily servings of high-residue produce had a 49% lower total sperm count and 32% fewer normally shaped sperm than men eating under half a serving — while total fruit and vegetable intake on its own was unrelated to semen quality.

Main groups of concern

Chemical group	Examples	Where exposure comes from	Hormone concern
Organochlorines (legacy)	DDT/DDE, dieldrin, PCBs (industrial cousins)	Banned in most countries decades ago but persist in fatty fish, meat, dairy, and in body fat	Estrogenic or anti-androgenic; linked to lower semen quality at high exposure
Organophosphates	Chlorpyrifos, malathion	Residues on conventional produce; farm and home pest control	Linked to poorer sperm quality and altered hormones
Triazine herbicides	Atrazine	Corn and sugarcane farming runoff; drinking water in farm regions	Disrupts the brain–pituitary signals that trigger ovulation; banned in the EU since 2004 but widely used in the US
Pyrethroids	Permethrin, cypermethrin	Household insect sprays, pet flea treatments, mosquito control	Anti-androgenic activity in lab studies; linked to lower sperm quality
Fungicides	Vinclozolin, mancozeb	Fruit, vegetables, vineyards	Anti-androgenic; vinclozolin is a classic lab model of male reproductive disruption
Glyphosate	Roundup and others	The most-used herbicide worldwide; grains, legumes	Human hormone evidence is limited and debated; lab studies suggest effects on steroid production

Who has higher risk

- Farmworkers, pesticide applicators, groundskeepers and their families
- People living next to heavily sprayed fields
- Private-well users in agricultural areas
- Pregnant women and young children (developing systems are most sensitive)

What to do

- Keep eating plenty of fruit and vegetables — the benefits are clear.

- Buy organic for the highest-residue crops if budget allows. In the EARTH study, high-residue items included peppers, spinach, strawberries, apples and pears; low-residue items included peas, beans, onions and grapefruit. The Environmental Working Group's yearly "Dirty Dozen" and "Clean Fifteen" lists help you prioritize. Find these at EWG.org
- Soak produce in a mixture of baking soda and lemon juice or vinegar; peel where sensible (such as with citrus). This lowers but doesn't entirely remove residues.
- Trim fat from meat and choose smaller, low-on-the-food-chain fish to reduce persistent organochlorines.
- Avoid home and garden pesticide sprays and "bug bombs"; use traps, screens and baits instead.
- Leave shoes at the door — pesticides track in on soles and build up in dust.
- If you work with pesticides: wear gloves and protective gear, change clothes before going home, and wash work clothes separately.

Chapter 6: Personal care and household chemicals

What you put on your skin each day is a major and very changeable source of exposure. In the HERMOSA study, 100 teenage girls who switched to products free of phthalates, parabens, triclosan and oxybenzone for just three days cut methyl and propyl parabens by about 44–45%, triclosan by 36%, oxybenzone by 36% and the fragrance phthalate DEP by 27%.

Women tend to carry more of these chemicals because they use more products: about 12 a day for adult women and 17 for teenage girls, according to one survey.

Chemical	What it does in products	Where it hides	Label words to watch
Parabens	Preservative	Lotions, makeup, shampoo, some foods and drugs	methylparaben, propylparaben, butylparaben
Phthalates (esp. DEP)	Makes scent last	Perfume, body spray, scented lotion, air fresheners, candles	"fragrance," "parfum" — rarely listed by name
Triclosan, triclocarban	Antibacterial	Older antibacterial soaps, some toothpastes, deodorants, "antimicrobial" textiles and cutting boards	triclosan, triclocarban, "Microban"
Oxybenzone (BP-3) and other UV filters	Absorbs UV light	Chemical sunscreens, moisturizers and lip balm with SPF	oxybenzone, benzophenone-3, octinoxate, homosalate
Cyclic siloxanes	Smooth, silky feel	Hair products, deodorants, foundations	cyclopentasiloxane (D5), cyclotetrasiloxane (D4)
Flame retardants	Slow burning	Older foam furniture, carpet padding, electronics, car seats	PBDEs, organophosphate flame retardants (rarely labeled)
Alkylphenols	Detergent/surfactant	Some industrial cleaners, paints, pesticides	nonylphenol, octylphenol, "nonoxynol"

How they act

- **Parabens** weakly mimic estrogen; longer-chain ones (propyl-, butyl-) are more potent.
- **Triclosan** can disrupt thyroid hormone and has been linked to lower ovarian reserve in some studies. The US FDA removed it from over-the-counter antibacterial hand soaps in 2016.
- **Oxybenzone** shows estrogen- and androgen-disrupting activity in lab studies.
- **Flame retardants (PBDEs)** interfere with thyroid hormone and have been linked to longer time to pregnancy.

What to do

- Cut the number of products you use daily — fewer products means fewer exposures.
- Choose “fragrance-free” (not “unscented,” which can contain masking fragrance).
- Skip antibacterial soaps; plain soap and water work just as well.
- Use mineral sunscreens (zinc oxide or titanium dioxide) when sun protection is needed.
- Replace plug-in air fresheners, scented candles and fabric softener sheets with essential oils, ventilation, unscented options or nothing.
- Use simple cleaners: plain soap, vinegar, baking soda, or certified “Safer Choice” products.
- When replacing a sofa or mattress, look for no added flame retardants (in California, the TB117-2013 label discloses this).
- Apps and databases such as EWG.org’s Skin Deep or Think Dirty can help you assess the toxic impact of different products, but they are guides, not definitive safety ratings.

Chapter 7: Heavy metals, air pollution and lifestyle toxins

Metals, polluted air, tobacco and alcohol harm fertility mainly through oxidative stress and hormone disruption — and smoking alone delivers several of them at once. A 2025 review concluded that metals such as arsenic, cadmium, chromium, mercury and lead reduce sperm quality in men and interfere with egg maturation and ovarian function in women, even at low levels.

Heavy metals

Metal	Main sources	Reproductive concerns
Lead	Old paint (pre-1978 homes), old pipes, some imported ceramics, spices and cosmetics, firing ranges, renovation dust	Lower sperm count and quality; stored in bone and released during pregnancy; miscarriage and fetal harm
Cadmium	Cigarette smoke (the biggest source), some rice, leafy greens and shellfish grown in polluted soil, batteries, pigments	Damages the testes' blood barrier; acts as a "metalloestrogen"; linked to poorer sperm and egg quality
Mercury	Large predatory fish (swordfish, shark, king mackerel, bigeye tuna), old dental amalgam, some skin-lightening creams	Linked to infertility and poorer semen quality; harmful to the developing fetal brain
Arsenic	Well water in some regions, rice and rice products	Oxidative stress, lower sperm motility; pregnancy complications

Air pollution

Fine particles (PM2.5) from traffic, industry and wildfire smoke are linked to lower ovarian reserve and poorer sperm. In a Boston fertility clinic study, every 2 µg/m³ increase in PM2.5 was associated with a 7.2% lower antral follicle count — a sign air pollution may speed up reproductive aging. Even in young egg donors, higher PM2.5 was linked to 6–8% fewer mature eggs.

Tobacco, vaping, alcohol and cannabis

- **Smoking** exposes you to cadmium, lead, polycyclic aromatic hydrocarbons and many other toxins. It speeds egg loss, can bring menopause earlier, and lowers sperm count, motility and DNA quality. Secondhand smoke matters too.
- **Vaping** is less studied, but e-liquids and aerosols contain nicotine, metals and flavoring chemicals; nicotine itself affects hormones.
- **Alcohol** at heavy levels lowers testosterone and disrupts ovulation. Experts advise no alcohol once you may be pregnant.

- **Cannabis** has been linked to lower sperm counts and disrupted ovulation in some studies.

Heat and the body (for men)

Heat isn't a chemical, but it compounds toxin effects on sperm: hot tubs, saunas, laptops on the lap and tight underwear can all temporarily lower sperm quality.

What to do

- **Stop smoking** — the single biggest reduction in toxic metal exposure for smokers. Ask your doctor about support; both partners benefit.
- Test your home for lead if built before 1978, and use lead-safe methods for renovations. Check imported pottery, spices and cosmetics.
- Choose low-mercury fish (salmon, sardines, trout, anchovies, shrimp, cod) 2–3 times a week; avoid shark, swordfish, king mackerel, tilefish, marlin and bigeye tuna.
- Cook rice in plenty of water (like pasta) and drain it to lower arsenic; vary your grains. Always wash rice well before cooking!
- Test private well water for arsenic, lead and nitrates.
- On bad air days, limit outdoor exercise near traffic; use a HEPA air purifier at home, especially during wildfire smoke.
- Keep alcohol low (or none while trying to conceive) and avoid cannabis.

Chapter 8: Effects on women's fertility

In women, endocrine disruptors are most consistently linked to a smaller egg supply, hormone conditions such as PCOS and endometriosis, and lower success with IVF. Because a woman is born with all the eggs she will ever have, damage to the ovarian reserve can't be replaced — which makes prevention especially valuable.

Outcome	Chemicals most often linked	What studies show
Lower ovarian reserve (AMH, antral follicle count)	Phthalates, BPA, PFAS, air pollution (PM2.5), smoking	Fewer follicles; earlier reproductive aging; cigarette smoking can bring menopause forward by 1–4 years
Ovulation and cycle problems	BPA, PFAS, atrazine, heavy metals	Irregular or longer cycles; disrupted LH surge
PCOS	BPA, PFAS	Women with PCOS often have higher blood levels; BPA may raise androgens and worsen insulin resistance
Endometriosis	Phthalates, PFAS, dioxins, PCBs	Associations with higher risk; estrogen-like chemicals may feed lesions
Egg quality and IVF results	BPA, phthalates, PFAS, PM2.5	Chemicals found in follicular fluid; fewer mature eggs, lower fertilization and implantation rates
Time to pregnancy	PFAS, phthalates, flame retardants	Longer time to conceive; in one PFAS study, 30–40% lower odds of pregnancy within a year
Pregnancy outcomes	Lead, mercury, PFAS, phthalates, air pollution	Miscarriage, preterm birth, low birth weight, pregnancy high blood pressure
Early menopause	Smoking, PFAS, phthalates, some pesticides	Earlier loss of ovarian function

Key points

- **The womb is a critical window.** A mother's exposures during pregnancy can affect her daughter's future egg supply — and her son's future sperm count (see Chapter 9).
- **The chemicals reach the egg.** BPA, phthalates and PFAS have all been measured in the follicular fluid that surrounds developing eggs.
- **Effects may add to other factors.** Age, weight, insulin resistance and stress all affect fertility. Obesity can increase the body's store of fat-soluble chemicals, and EDCs may themselves promote weight gain ("obesogens").
- **Many links come from fertility-clinic patients.** These women are more likely to have existing problems, so results may not apply equally to everyone — but they are most relevant to people trying to conceive.

Signs worth discussing with a doctor

- Cycles shorter than 21 or longer than 35 days, or very irregular
- No periods for 3+ months (when not pregnant)
- Signs of high androgens: acne, excess facial or body hair, scalp hair thinning
- Very painful periods or pain with sex (possible endometriosis)
- Not pregnant after 12 months of trying (6 months if 35+)
- Two or more miscarriages

Chapter 9: Effects on men's fertility

Sperm counts have fallen by about half in under 50 years, and anti-androgenic chemicals are a leading suspect. A 2022 meta-analysis of 57,000+ men across 53 countries found average sperm concentration dropped from about 101 million/mL to 49 million/mL between 1973 and 2018, with the decline speeding up from 1.2% a year to 2.6% a year after 2000.

That study didn't test causes, and some experts argue older counting methods inflate the early numbers. The average today is still within the WHO normal range (above 15 million/mL). But the decline tracks other worrying trends — rising testicular cancer, falling testosterone and more genital birth defects — that point to hormone disruption, especially before birth.

Outcome	Chemicals most often linked	What studies show
Lower sperm count and concentration	Phthalates (DEHP, DBP), PFAS, pesticides, PCBs, lead, cadmium, air pollution, smoking	Consistent associations across many studies; high pesticide-residue produce linked to 49% lower total count in one cohort
Poor motility and shape	Phthalates, PFAS, organophosphates, pyrethroids, heavy metals	Fewer normally formed and fast-swimming sperm
Sperm DNA damage	Phthalates, BPA, PAHs from smoke and pollution, metals	Higher DNA fragmentation, linked to lower fertilization rates and miscarriage
Lower testosterone	Phthalates, PFAS, BPA, cadmium, some pesticides	Phthalates block testosterone production in Leydig cells
Birth defects in sons	Mother's phthalate exposure in pregnancy	Shorter anogenital distance, undescended testes, hypospadias — together called testicular dysgenesis syndrome

Key points

- **Two windows matter.** A man's fertility depends both on his mother's exposures while he was in the womb and on his own exposures now.
- **The 3-month rule.** Sperm take about 70–90 days to develop. Changes you make today — quitting smoking, cutting phthalates, avoiding heat — show up in a semen test about three months later. That makes men's fertility unusually responsive to lifestyle change.
- **Men's exposures affect pregnancy too.** Sperm DNA damage can affect embryo quality and miscarriage risk, so preconception care is for both partners.
- **Occupation matters.** Farmers, pesticide applicators, painters, welders, firefighters and plastics or battery workers can have much higher exposures.

Signs worth discussing with a doctor

- Not conceiving after 12 months (6 if your partner is 35+) — a semen analysis is simple and should happen early, alongside testing for her
- Low libido, erectile problems, fatigue or loss of muscle (possible low testosterone)
- Testicular pain, swelling or a lump
- History of undescended testes, chemotherapy or known high chemical exposure at work

Chapter 10: Your practical action plan

Start with the few changes that cut the most exposure for the least effort, ideally at least three months before trying to conceive — both partners together. Perfection isn't the goal; steady reduction is.

Phase 1: The big five (this week)

1. **Stop smoking and avoid secondhand smoke.** The largest single toxic load for smokers.
2. **Stop heating food in plastic.** Microwave and store hot food in glass or ceramic.
3. **Swap canned food for fresh, frozen, dried or glass-jarred.** Cuts BPA quickly.
4. **Go fragrance-free.** Lotion, laundry detergent, air fresheners, candles — the main hidden phthalate source.
5. **Check your water.** Read your utility report or test your well; filter if PFAS, lead or arsenic show up.

Phase 2: Kitchen and food (this month)

- Cook from fresh ingredients more often; cut fast food and takeout.
- Buy organic for high-residue produce when you can; wash all produce well.
- Replace scratched non-stick pans with stainless steel, cast iron or ceramic.
- Store leftovers in glass or stainless steel; use beeswax wraps or parchment instead of cling film on hot or fatty foods.
- Choose low-mercury fish; trim fat from meat; eat a varied diet.
- Avoid microwave popcorn bags and greasy paper wrappers.

Phase 3: Bathroom and personal care (next 1–3 months)

- Replace products as they run out — no need to throw everything away at once.
- Choose paraben-free, triclosan-free and fragrance-free options.
- Switch to mineral sunscreen.
- Simplify: fewer products, fewer exposures.
- For nail polish and hair dyes, use less often and in well-ventilated spaces.

Phase 4: Home environment (ongoing)

- Shoes off at the door.

- Wet-dust and HEPA-vacuum weekly — dust carries flame retardants, phthalates, PFAS and pesticides.
- Open windows daily for fresh air (unless outdoor air quality is poor); use a HEPA air purifier during smoke events.
- Avoid pesticide sprays indoors and on lawns.
- Choose furniture, mattresses and carpets without stain-guard or added flame retardants when replacing them.
- If your home was built before 1978, test for lead before renovating.

Phase 5: Work

- Know what chemicals you handle; ask for safety data sheets.
- Use the gloves, masks and ventilation provided.
- Change out of work clothes and shoes before going home; wash them separately.
- If you're planning a pregnancy, talk to your employer or occupational health about reducing exposure.

Next: support your body's own elimination pathways (Chapter 11)

Chapter 11: Lowering your toxic burden naturally

Your body already runs a sophisticated elimination system every day; nutrition, movement, hydration, sleep and calm keep it working well. Cutting exposure (Chapter 10) always comes first — these five pillars then help your body clear what still gets in. Some of the evidence comes from clinical trials, some is early or from animal studies; each section says which.

Your elimination pathways

Organ	What it does	Chemicals it handles	How to support it
Liver	Phase I enzymes (CYP450) alter chemicals; phase II attaches a molecule (glucuronide, sulfate, glutathione, methyl group) to make them water-soluble. Also breaks down used estrogen	BPA, phthalates, parabens, pesticides, hormones	Cruciferous vegetables, adequate protein, B vitamins, little or no alcohol
Gut	Bile carries processed chemicals and spent estrogens into the intestine for removal in stool. Some gut bacteria can undo this work and let them be reabsorbed	Estrogens, PCBs, dioxins, some metals	Fiber, fermented foods, plants for a diverse microbiome, daily bowel movements
Kidneys	Filter water-soluble metabolites into urine	Most BPA, phthalate, paraben and pesticide metabolites	Good hydration
Skin	Releases some compounds in sweat	Some metals, BPA, phthalates (not PFAS)	Regular sweat-inducing exercise
Lungs	Exhale volatile compounds	Solvents and other volatile chemicals	Clean indoor air, aerobic activity

1. Nutrition: feed your detox pathways

Eat cruciferous vegetables daily. Broccoli, broccoli sprouts, kale, cabbage, Brussels sprouts, cauliflower, arugula and watercress contain glucoraphanin, which becomes sulforaphane — a strong activator of phase II enzymes and glutathione. In a 12-week trial of 291 people in a polluted region of China, a daily broccoli-sprout drink increased urinary excretion of benzene by 61% and acrolein by 23% compared with placebo. Crucifers also contain compounds (indole-3-carbinol, DIM) involved in healthy estrogen metabolism.

- *Tip:* chop or chew crucifers well and let chopped broccoli sit for a few minutes before cooking; lightly steam rather than boil; sprinkle mustard seed powder on cooked crucifers to restore the enzyme that makes sulforaphane.

Get 25–38 g of fiber a day. Fiber binds bile-excreted chemicals and spent estrogens in the gut so they leave in stool rather than being reabsorbed. Human evidence is limited but supportive: in a clinical trial in patients poisoned with PCB-contaminated rice oil, rice bran fiber plus cholestyramine increased PCB excretion, and animal studies show various fibers increase fecal excretion of PCBs and dioxins. Good sources: ground flaxseed, chia, oats, legumes, vegetables, berries, whole grains.

Eat enough protein. Phase II detoxification uses amino acids — glycine, cysteine, taurine, glutamine. Include legumes, eggs, fish, poultry, nuts and seeds at each meal.

Build glutathione, your master antioxidant. Sulfur-rich foods (garlic, onions, leeks, eggs, crucifers), selenium (1–2 Brazil nuts a day — no more), and vitamin C (citrus, kiwi, peppers) all support it.

Support methylation. Folate (leafy greens, legumes), B12 (animal foods or supplement), B6 and choline (eggs) are needed for clearing hormones and chemicals — and for a healthy pregnancy.

Use minerals to block toxic metals. Toxic metals compete with essential minerals for absorption, so a diet rich in calcium, iron and zinc absorbs less lead and cadmium, and selenium binds mercury. In a randomized trial of 670 pregnant women in Mexico City, 1,200 mg of daily calcium lowered blood lead by 11% on average, and by 24% in women who took it consistently. Avoid bone-meal or dolomite calcium, which can contain lead.

Protect eggs and sperm with antioxidants. Oxidative stress is a main way toxins damage eggs, sperm and sperm DNA. Eat a rainbow of vegetables and fruit, nuts, seeds, olive oil and oily low-mercury fish (omega-3s). Supplements such as CoQ10, vitamin E, zinc and selenium are widely used in naturopathic fertility care; trial evidence is mixed, so use food first and discuss supplements with your practitioner.

Lighten the liver's load. Limit alcohol (it competes for liver capacity and depletes glutathione), added sugar and ultra-processed foods.

About liver herbs. Herbs such as milk thistle, dandelion root and turmeric are traditional naturopathic liver supports. They are mainly studied in liver disease rather than chemical clearance, and some are not advised when trying to conceive or during pregnancy — use them only with a qualified practitioner.

2. Physical activity: move and sweat

Why it helps

- Builds muscle and improves insulin sensitivity, which helps balance hormones in PCOS and supports testosterone in men.
- Reduces body fat, the main storage site for persistent chemicals.
- Boosts circulation, lymph flow and bowel regularity.

Sweat as an exit route. Small studies by Genuis and colleagues found BPA in the sweat of 16 of 20 people, including some with none detectable in blood or urine, and some phthalates and metals in sweat too. However, PFAS were not excreted efficiently in sweat. This is promising but early evidence.

A caution on heat. Saunas are a traditional naturopathic tool, but heat temporarily lowers sperm quality — men trying to conceive should skip saunas and hot tubs and sweat through exercise instead. Women who may be pregnant should avoid overheating.

Lose weight gradually. Fat stores persistent chemicals, and losing it releases them: a review found blood levels of most persistent pollutants rose 2–4% per kilogram lost, staying elevated a year later. The benefits of reaching a healthy weight still outweigh this, but aim for slow, steady loss with a high-fiber, plant-rich diet, ideally well before trying to conceive.

How much: about 150 minutes of moderate activity a week plus two strength sessions. Very intense training can suppress ovulation and sperm production — balance is the aim.

3. Hydration: keep the kidneys flowing

- Your kidneys clear most metabolites of BPA, phthalates, parabens and pesticides in urine. Steady fluid intake keeps that route open.
- Aim for pale-yellow urine. As a guide, total daily fluid from all drinks and food is about 2.7 L for women and 3.7 L for men, more in heat or with exercise.
- Make your water clean: filter tap water if testing shows contaminants (Chapter 4), and carry it in glass or stainless steel rather than plastic.
- Herbal teas such as ginger, peppermint or rooibos count toward fluids; check pregnancy safety for others.
- Keep caffeine moderate (under about 200 mg a day when trying to conceive) and avoid sugary drinks.

4. Sleep: the nightly reset

- Deep sleep is when the body repairs tissue and the brain's waste-clearing (glymphatic) system is most active. This is especially true during the period between 10 PM and 2 AM, so sleep by 10 PM is important!
- Reproductive hormones follow sleep rhythms: testosterone rises during sleep, and melatonin acts as an antioxidant in the fluid around developing eggs.
- Short or disrupted sleep raises cortisol and insulin resistance and is linked to lower testosterone; shift work is linked to menstrual irregularity.

Habits that help

- 7–9 hours a night on a consistent schedule
- A dark, cool, quiet bedroom; screens off an hour before bed
- Morning daylight to set your body clock
- Little caffeine after midday and little alcohol in the evening

5. Stress management: calm the hormone axis

- Chronic stress raises cortisol, which can suppress the brain's fertility signals (GnRH, LH, FSH) — leading to irregular ovulation, lower testosterone and poorer sperm.
- Stress and chemical toxins act through the same routes: oxidative stress and inflammation.
- Stress also drives poor sleep, comfort eating and alcohol — undermining every other pillar.
- Trying to conceive is itself stressful, for both partners.

Practices that help

- Daily mind–body practice: breathwork, meditation, yoga or tai chi (even 10 minutes)
- Time in nature and gentle outdoor movement
- Connection: friends, partner time, support groups, counseling
- Acupuncture and adaptogenic herbs (such as ashwagandha) are common in naturopathic fertility care; evidence is mixed, and some herbs are not suitable in pregnancy — use them under professional guidance.

Approaches to use with care or avoid

- **Juice cleanses and prolonged fasts** can mobilize stored chemicals quickly while starving the liver of the protein it needs for phase II detoxification.
- **Unsupervised chelation, "heavy metal detox" kits, colonics and detox foot pads** are unproven and some are harmful.
- **Herbal detox supplements** while trying to conceive or pregnant should only be used with a qualified practitioner.
- **An evidence-based option for high PFAS:** in a randomized trial of 285 firefighters, donating plasma every 6 weeks lowered PFAS by about 30% over a year, and blood every 12 weeks by about 10%. This may suit people with known high levels who are eligible donors — but not during pregnancy; discuss it with your practitioner.

A sample supportive day

- **Morning:** a glass of filtered water; 10 minutes of daylight and gentle movement; breakfast of oats with ground flaxseed, berries and nuts.
- **Midday:** lunch with a big serving of leafy or cruciferous vegetables, legumes or fish, and olive oil; a brisk 20–30 minute walk.
- **Afternoon:** refill your water bottle; herbal tea instead of a second coffee.
- **Evening:** dinner with colorful vegetables and protein; screens off an hour before bed; 10 minutes of breathwork or stretching; lights out for 7–9 hours.

Chapter 12: Testing, your practitioners, and keeping perspective

Standard fertility tests come first; chemical testing is optional and mostly useful for persistent chemicals or known high exposure. A normal fertility workup tells you far more about your chances than a toxin panel.

Standard fertility tests to ask about

For	Test	What it shows
Women	AMH and antral follicle count (ultrasound)	Ovarian reserve (egg supply)
Women	Day 2–5 FSH, LH, estradiol	Ovarian and pituitary function
Women	Mid-luteal progesterone	Whether ovulation happened
Women	TSH, prolactin, testosterone, glucose/insulin	Thyroid, pituitary and PCOS screening
Women	Tubal test (HSG or HyCoSy)	Whether fallopian tubes are open
Men	Semen analysis (repeat if abnormal)	Count, motility, shape, volume
Men	Testosterone, LH, FSH	Hormone production
Men	Sperm DNA fragmentation (in selected cases)	DNA damage, e.g. after repeated IVF failure or miscarriage
Both	Blood lead	Useful if older housing, certain jobs or hobbies

Chemical (biomonitoring) testing

- **PFAS blood test.** The US National Academies recommend PFAS testing for people with likely elevated exposure (for example, contaminated water or occupational exposure). Results help guide care and future exposure reduction.
- **Blood lead and mercury.** Widely available and meaningful.
- **Urine BPA, phthalates and parabens.** These reflect only the last day or two, so a single result says little about long-term exposure. Better used before and after changes, or in research.
- **Be cautious with direct-to-consumer "toxin panels"** and hair analysis — methods vary, and results are often hard to interpret.

Working with your naturopathic doctor and medical team

- Consider a naturopathic doctor (ND) to coordinate diet, lifestyle, herbs and supplements safely alongside your GP, gynecologist, urologist or fertility specialist — and make sure each knows

what the other recommends. Bring a short exposure history: your water source, job, hobbies, home age, smoking, fish intake and product use.

- Ask: “Could environmental exposures be relevant for us, and is any testing worthwhile?”
- Ask for a semen analysis early — male factor is involved in about half of couples’ infertility.
- Organizations such as ACOG (American College of Obstetricians and Gynecologists) and FIGO (International Federation of Gynecology and Obstetrics) recommend that clinicians counsel patients about reducing toxic environmental exposures, so this is a legitimate question.

Keeping perspective

- **Exposure is universal; harm depends on dose, timing and mixtures.** Nearly everyone carries some of these chemicals. Having them doesn’t mean you will be infertile.
- **Small, steady changes work.** Short-lived chemicals drop within days of reducing exposure.
- **Don’t let worry crowd out the basics.** Age, timing intercourse in the fertile window, a healthy weight, not smoking and treating medical conditions remain the biggest factors you control.
- **Stress and guilt don’t help.** You cannot remove every chemical. Focus on the big wins and let the rest go.
- **Advocate as well as adapt.** Personal choices help, but the largest reductions come from regulation — supporting stronger chemical safety rules protects everyone, including those who can’t afford organic food or water filters.

Quick reference

At-a-glance chemical guide

Chemical	Stays in body	Top sources	Best single swap
BPA / bisphenols	Hours	Can linings, hard plastics, receipts	Fresh or frozen instead of canned
Phthalates	Hours–days	Fragrance, vinyl, processed and fast food	Fragrance-free products; cook at home
Parabens	Hours–days	Lotions, makeup, shampoo	Paraben-free personal care
Triclosan	Days	Antibacterial soaps and products	Plain soap
Oxybenzone	Days	Chemical sunscreen	Mineral sunscreen
PFAS	Years	Water, grease-proof packaging, non-stick pans	Certified water filter; stainless or cast iron pans
Pesticides (current-use)	Days	Conventional high-residue produce, home sprays	Organic for high-residue crops; no home sprays
Legacy organochlorines, PCBs	Years	Fatty animal foods, contaminated fish	Trim fat; choose smaller fish
Lead	Years (bone)	Old paint, pipes, imported goods	Test home and water
Mercury	Weeks–months	Large predatory fish	Low-mercury fish
Cadmium	Decades	Cigarette smoke	Quit smoking

Preconception checklist (both partners, about 3 months ahead)

- ☐ Stop smoking and vaping; avoid secondhand smoke
- ☐ Limit or stop alcohol; avoid cannabis
- ☐ Check water quality; install a certified filter if needed
- ☐ No more heating food in plastic
- ☐ Replace canned food with fresh, frozen or jarred
- ☐ Switch to fragrance-free personal care, laundry and cleaning products
- ☐ Replace scratched non-stick cookware
- ☐ Buy organic for highest-residue produce where possible

- ☐ Choose low-mercury fish
- ☐ Shoes off at the door; damp-dust and HEPA-vacuum weekly
- ☐ Review workplace exposures and protective gear
- ☐ Book a preconception check-up; ask about semen analysis and relevant tests
- ☐ Women: start folic acid as advised
- ☐ Eat cruciferous vegetables daily and 25–38 g of fiber
- ☐ Drink enough filtered water for pale-yellow urine
- ☐ Move 150+ minutes a week, with strength work twice
- ☐ Sleep 7–9 hours on a regular schedule
- ☐ Practice a daily stress-reduction habit
- ☐ Men: skip saunas, hot tubs and laptops on the lap

Glossary

- **AMH (anti-Müllerian hormone):** blood marker of remaining egg supply.
- **Anogenital distance:** distance from anus to genitals; shorter in boys is a marker of lower testosterone exposure before birth.
- **Anti-androgenic:** blocking or lowering the action of male hormones such as testosterone.
- **Antral follicle count:** number of small egg-containing follicles seen on ultrasound.
- **EDC (endocrine-disrupting chemical):** an outside chemical that interferes with hormones.
- **Estrogenic:** mimicking the female hormone estrogen.
- **Fecundability:** the chance of conceiving in a single menstrual cycle.
- **Glutathione:** the body's main antioxidant, used by the liver to neutralize and package toxins.
- **Half-life:** time for the body to clear half of a chemical.
- **HPG axis:** the hypothalamus–pituitary–gonad hormone chain that controls fertility.
- **Naturopathic doctor (ND):** a practitioner trained in natural therapies, nutrition and lifestyle medicine; licensing varies by country and state.
- **Obesogen:** a chemical that may promote fat gain by disrupting metabolism.
- **PCOS (polycystic ovary syndrome):** a common hormone disorder causing irregular ovulation and high androgens.
- **PFAS:** per- and polyfluoroalkyl substances, or "forever chemicals."
- **Phase I / phase II detoxification:** the liver's two-step process of altering chemicals and then attaching a molecule so they can be excreted.
- **PM2.5:** fine particles in air pollution, 2.5 micrometers or smaller.
- **Sulforaphane:** a compound from broccoli and other crucifers that switches on detoxification enzymes.
- **Testicular dysgenesis syndrome:** a cluster of male reproductive problems (low sperm, testicular cancer, genital defects) thought to start before birth.

Sources

- Tzouma Z, et al. Associations Between EDC Exposure and Fertility Outcomes: A Decade of Human Epidemiological Evidence. *Life*, 2025.
- Bisphenol A and DEHP: Hidden threats to female fertility. *J Obstet Gynaecol*, 2025.
- Cohen NJ, et al. Exposure to PFAS and women's fertility outcomes in a Singaporean preconception cohort. *Sci Total Environ*, 2023.
- Joensen UN, et al. Do perfluoroalkyl compounds impair human semen quality? *Environ Health Perspect*, 2009.
- Radke EG, et al. Phthalate exposure and male reproductive outcomes: a systematic review. *Environ Int*, 2018.
- Rudel RA, et al. Food packaging and BPA and DEHP exposure: findings from a dietary intervention. *Environ Health Perspect*, 2011.
- Harley KG, et al. The HERMOSA intervention study. *Environ Health Perspect*, 2016; study background.
- Chiu YH, et al. Fruit and vegetable intake and their pesticide residues in relation to semen quality. *Hum Reprod*, 2015.
- Wirbisky SE, Freeman JL. Atrazine exposure and reproductive dysfunction through the HPG axis. *Toxics*, 2015.
- Levine H, et al. Temporal trends in sperm count (*Hum Reprod Update*, 2022), as reported by Down To Earth and Gigazine.
- Gaskins AJ, et al. Exposure to fine particulate matter and ovarian reserve. *Epidemiology*, 2019.
- LaPointe S, et al. Traffic-related air pollution and oocyte quality in egg donors. *Environ Int*, 2024.
- Marconi G, et al. Heavy metals and human reproductive toxicity. *Reprod Toxicol*, 2025.
- European Commission. EU prohibition on BPA in food contact materials from 20 January 2025.
- Harvard Environmental & Energy Law Program. PFAS in drinking water regulatory tracker (updated May 2026).
- Egner PA, et al. Rapid and sustainable detoxication of airborne pollutants by broccoli sprout beverage. *Cancer Prev Res*, 2014.
- Rice bran fiber and cholestyramine for PCB excretion in Yu-Cheng patients.
- Ettinger AS, et al. Effect of calcium supplementation on blood lead levels in pregnancy. *Environ Health Perspect*, 2009.

- Genuis SJ, et al. Human excretion of bisphenol A: Blood, Urine, and Sweat (BUS) study, 2012; PFCs and PCBs through perspiration, 2013.
- Increased blood levels of persistent organic pollutants in obese individuals after weight loss: a review. *J Toxicol Environ Health B*, 2017.
- Effect of plasma and blood donations on PFAS levels in firefighters: a randomized clinical trial. *JAMA Netw Open*, 2022; plain-language summary.