

High-Risk Conditions	Assessment of Vulnerability	Maximum	High	Intermediate	Daily
Pregnancy	Third trimester of pregnancy	Telecommuting from home	Telecommuting from home	May return to non-COVID* work-related areas	Full return
Breastfeeding	During maternity leave	Telecommuting from home	Telecommuting from home	Telecommuting from home	Full return
Obesity	BMI $\geq 40$ kg/m <sup>2</sup>	Telecommuting from home	Telecommuting from home	May return to non-COVID* work-related areas	Full return
People over 60 without disorders	Over 60 without a disorder	Telecommuting from home	Telecommuting from home	May return to non-COVID* work-related areas	Full return
Diabetes Mellitus	HbA1c > 8%	Telecommuting from home	Telecommuting from home	May return to non-COVID* work-related areas	Full return
Systemic Arterial Hypertension (SAH) uncontrolled	Stage 2 Hypertension (Systolic 160-179 mmHg or Diastolic 100-109 mmHg) or greater	Telecommuting from home	Telecommuting from home	May return to non-COVID* work-related areas	Full return
Chronic Obstructive Pulmonary Disease (COPD)	Diagnosis established by Pneumology. Requires daily treatment with CAT $\geq 10$	Telecommuting from home	Telecommuting from home	May return to non-COVID* work-related areas	Full return
Asthma	Moderate. (GINA) b Diagnosis established by Pneumology. Has daily symptoms that affect activity and sleep. Requires quick relief mediation daily	Telecommuting from home	Telecommuting from home	May return to non-COVID* work-related areas	Full return
Cardiovascular or Cerebrovascular Diseases	Diagnosis established by cardiology or neurology. Continued treatment required or has required hospitalization because of pathology over the past year, not including arterial hypertension	Telecommuting from home	Telecommuting from home	May return to non-COVID* work-related areas	Full return
Chronic Kidney Disease	Workers in stage 3b KDIGO or higher, with proteinuria events in the last year, or some other manifestation of renal involvement	Telecommuting from home	Telecommuting from home	May return to non-COVID* work-related areas	Full return
Immunosuppressive treatment in pathologic stages	Pathologic stages that require immunosuppressive treatment	Telecommuting from home	Telecommuting from home	May return to non-COVID* work-related areas	Full return
HIV infection	CD4 < 350 cells/mL	Telecommuting from home	Telecommuting from home	May return to non-COVID* work-related areas	Full return
Cancer	With chemotherapeutic treatment in the last month	Telecommuting from home	Telecommuting from home	Telecommuting from home	Full return
Combination of concurrent disorders	Two or more disorders that meet the criteria for the assessment of vulnerability	Telecommuting from home	Telecommuting from home	Telecommuting from home	Full return
Combination of concurrent disorders	Two or more disorders that do not meet the criteria for the assessment of vulnerability	Full return	Full return	Full return	Full return