

Purpose: Provide proper patient screening for tobacco and smokeless tobacco use

Tobacco Screening:

- Tobacco Screening BPA fires in Rooming Navigator if patient has not been screened for tobacco use or smokeless tobacco use in past year.
- Any clinical staff can screen patients for tobacco use.
- Clear the Screening BPA by using one of the blue links in the screening BPA.
- Complete **BOTH** 'Tobacco Use' and 'Smokeless Tobacco Use' fields using options in examples below.
- Click "Mark as Reviewed" when done.

BestPractice Advisories 

Tobacco Screening BPA

Moderate (2)

! **Tobacco Use Screening Recommended**



Patient has not been screened for smoking and/or smokeless tobacco use in the past year.

Assess Smoking and Smokeless Tobacco Use then **Mark as Reviewed**

[Open Vitals Section \(Rooming Tab\)](#) 

[Open Vitals Section \(Visit Navigator\)](#) 

MA Screening

Available tobacco use selections when screening patients

Tobacco Use:  

Start Date:

Category Select

Search:

▲ Title
Current Every Day Smoker
Current Some Day Smoker
Former Smoker
Heavy Tobacco Smoker
Light Tobacco Smoker
Never Assessed
Never Smoker
Passive Smoke Exposure - Never Smoker
Smoker, Current Status Unknown
Unknown If Ever Smoked

10 categories loaded.

 **Accept**  **Cancel**

EXAMPLES: Patient Answers & Associated Tobacco Use Selections

- Use "**Never Smoker**" if patient uses **ONLY e-Cigarettes, e-Hoochak, or Vape Pens** and never used tobacco.
- Use "**Former Smoker**" if recently **switched** from regular cigarettes to e-cigs.
- Use "**Current Smoker**" when patient switches between cigarettes and e-cigs.
- Do **NOT** use "**Never Assessed**" or "**Unknown**".

Purpose: Complete Best Practice Advisory (BPA) for Tobacco Screening and Tobacco Counseling

Smoking is the most preventable cause of death in the US. It's linked to about one third of all deaths from health disease and 90% of lung cancers. ([AHA](#))

Basic Tobacco Counseling BPA for PCCs & MAs

Reason for Visit Device Select Vital Signs Extended Vitals **BestPractice**

Moderate (1)

ⓘ Tobacco Use Basic Counseling Recommended [Collapse](#) [A](#)

Smoking and/or smokeless tobacco use status indicates patient is a current tobacco user and basic counseling has not been performed in the past year.

Complete basic counseling then **Mark as Reviewed**

[Open Vitals Section \(Rooming Tab\)](#) [Open Vitals Section \(Visit Navigator\)](#)

- The Tobacco Counseling BPA will fire for patients identified as a tobacco user or smokeless tobacco user **every 3 months in specialty care clinics and every time a patient is identified as a tobacco user in primary care clinics or when counseling has not been done in the past year (whichever comes first).**
- To clear the Tobacco Counseling BPA:
 1. Click on the preferred blue link in BPA to Open Vitals Section (Rooming Tab or Visit Navigator)
 2. **Perform & document basic counseling** by using the scripting in the diagram below, then click **Mark as Reviewed** for **both patient interested or not in quitting.**
 3. Conclude basic counseling with **warm handoff** to provider

